

FORMATION AND CHARACTERIZATION OF SLOVAK DAIRY COMPOSITE SHEEP BREED: DESCRIPTION OF THE PROCESS: A REVIEW

M. MARGETÍN^{1,2*}, M. ORAVCOVÁ¹, J. HUBA¹, M. JANÍČEK²

¹NPPC – Research Institute for Animal Production Nitra, Slovak Republic

²Slovak University of Agriculture in Nitra, Slovak Republic

ABSTRACT

In the beginning of 1990s, the programme of formation of synthetic population of Slovak Dairy sheep (SD) was launched. Some flocks were involved into performance testing; the establishment of the breed was divided into two periods: first, crosses of local breeds (Improved Valachian, Tsigai and Merino) with improving breeds (Lacaune, East Friesian) of various proportions were formed; second, a close (*inter se*) breeding scheme within the crossbred population was applied. Parents of the next generation were chosen on the basis of breeding values for milk yield and litter size. At present, five thousands female SD individuals are included in the performance testing; 35 % yearling females enter the breeding scheme each year; a total, 50 thousands SD ewes are kept in commercial flocks. Milk and reproduction traits of SD were analyzed using mixed model methodology that included fixed and random effects. In similar way growth traits were analyzed. Milk traits analyses included 20,511 milk records belonging to 11,026 ewes kept in 52 flocks during the period between 1995 and 2015; reproduction traits analyses included 30,034 litter size records belonging to 9,671 ewes kept in 26 flocks during the period between 1997 and 2015. The standardized milk yield (SMY) and average daily milk yield (ADMY) increased more than twice: 79.8 ± 5.36 l (1995) vs. 164.0 ± 2.04 l (2015) and 495.1 ± 33.5 ml (1995) vs. 1035.3 ± 12.8 ml (2015), respectively. In ten percent of the most productive ewes, SMY was equal to 257.6 l, ADMY was equal to 1604 ml and dry matter was equal to 26.6 kg. Fat and protein contents decreased from 7.82 ± 0.101 % (1995) to 7.27 ± 0.040 % (2015) and from 5.83 ± 0.043 % (1995) to 5.69 ± 0.017 % (2015). Ewes of SD have good udder traits. Litter size in top 50 % flocks was above 150 %. Average daily gains from birth to weaning were 0.26 kg (female lambs) and 0.30 kg (male lambs). The population of SD was recognized as a breed in 2017.

Key words: dairy ewes; composite breed; milk traits; reproduction; growth

STATE OF THE ART

Following breeder's goals oriented towards high production, reproduction and functional traits of dairy sheep in Slovakia, the improving programme with intention of forming improved sheep population fitting semi-extensive production system was launched in Slovakia in 1990s. This was programme of forming Slovak Dairy sheep (SD). The breeding scheme was divided into two periods (Margetín and Čapistrák 1994; Margetín *et al.*, 2000 a,b,c). First, crosses of local breeds (Improved Valachian (IV), Tsigai (TS) and Merino

(M) with improving specialized breeds (Lacaune (LC), East Friesian (EF) of various proportions were formed in some flocks involved into performance testing. Preliminary evaluations of crossbreds with proportion of LC and EF ranged from 12.5 to 87.5 % were aimed at analyses of milk traits and litter size that are considered the most economically important traits (Apolen *et al.*, 2000; Čapistrák *et al.* 2000, 2002, 2005; Margetín *et al.*, 1993, 1999 a,b). Further analyses showed that the population with good udder shape, of appropriate cistern size and milkability was formed (Čapistrák *et al.*, 2006 a,b; Mačuhová *et al.*, 2008; 2009; Margetín *et al.*, 2005

*Correspondence: E-mail: margetin@vuzv.sk
Milan Margetín, NPPC – Research Institute for Animal Production Nitra,
Hlohovecká 2, 951 41 Lužianky, Slovak Republic
Tel.: +421 37 6546 313

Received: September 28, 2017

Accepted: October 2, 2017

b,c; Margetín *et al.*, 2011 a,b; Milerski *et al.*, 2005, 2006; Tančin *et al.*, 2011). Crossbreds with various proportions of LC and EF were also confirmed to be of good growth traits both till and after weaning (Margetín *et al.*, 2004 a,b). In addition, genetic parameters and variance components of udder morphology and milkability traits were preliminarily studied (Margetín *et al.* 2005a, 2008).

Second, a close (*inter se*), breeding scheme within the crossbred population was proposed (Margetín *et al.*, 2010, 2011c, 2012b) and applied taking into account results of analyses during the first period of the programme. The aim was to stabilize composite population using *inter se* (i.e. purebred) mating scheme that included dam and sire animals used as parents of the next generation that were selected according to their breeding values for milk and litter size traits. During this period, a lot of analyses were also done. Performance testing data of flocks involved in the programme were used for evaluation. These included analyses of udder morphology and milkability traits, since one goal was to form population of ewes that fit machine milking (Margetín *et al.*, 2013b), have good udder morphology and functional traits (Margetín *et al.*, 2012a; Makovický *et al.*, 2013, 2014, 2015a, 2017a), are of appropriate cistern size (Margetín *et al.*, 2011d; Makovický *et al.*, 2015b,c) and of minimal mastitis occurrence (Margetín *et al.*, 2013a). Growth curves of lambs of various proportions of LC and EF within IV and TS genotypes were compared (Makovický *et al.*, 2017b).

In 2016, a complex analysis of SD synthetic population involving period from 1995 to 2015 was done. Mixed model methodology (SAS, ver. 9.2) was employed. When milk yield during milking period (MY), standardized (SMY) and daily milk yield (DMY) as well as fat % (F %), protein % (P %), dry matter % (DM %), free of fat dry matter (DM-F %) and usable dry matter (UDM; kg) were investigated, following fixed effects: flock (52 levels), year (21 levels), parity (3 levels: first, second and third+ parity), litter size (4 levels: one lamb, two, three + lambs born and missing information about number of lambs born), covariates: milking period (in days) and interval between lambing and first test-day measurement (days) and random effect of ewe were considered. A total, 20,511 records that belonged to 11,026 ewes (1.86 records per ewe) entered the model for analyses of MY, SMY, DMY and 18,081 records entered the model for analyses of F %, P %, DM-F % and UDM. When litter size was investigated, the following fixed effects: flock (26 levels), year (19 levels) and age of dam (8 levels, one year old up to eight+ years old) and random effect of ewe were considered. A total, 30,034 litter size records that belonged to 9,671 ewes entered the model.

Study of Margetín (2016) confirmed that improving programme that resulted in acknowledgement of SD sheep with more favourable traits than those of local breeds was fulfilled. During the period of the last 21 years (from 1995 to 2015), SMY increased about twice (from 79.8 l in 1995 to 167.0 l in 2015). Similarly, DMY and production of usable dry matter increased by 109 and 79 %. Most productive flocks reached MY ranging from 170 to 190 l. Top 10 % ewes reached SMY above 250 l, DMY reached more than 1,600 ml and production of usable dry matter was above 26 kg. Average F % and P % slightly decreased (7 and 2.4 %, respectively). Also, ewes of SD population were of good udder morphology and functional udder traits. Prolificacy did not fulfill expectations and was 141.1 %. Nevertheless, it was about 160 % in top flocks. Average daily gain till weaning was about 300 g in male lambs and about 260 g in female lambs. The frequency of ARR allele of the prion gene ranged from 0.711 to 0.789 (Margetín *et al.*, 2016).

Results of SD sheep flocks resulted in fact that this population was recognized as a native breed on May 26, 2017. Detailed information about history of breeding this population can be found in the study of Margetín *et al.* (2017). Moreover, the importance of development of this population from social and breeders' point of view is described in the reports of Margetín (2017a,b).

CONCLUSION

Population of Slovak Dairy sheep is fairly large (more than 5 ths. females in performance testing, 35 % yearling females enter mating schemes), so it can be successfully bred in Slovakia in the future. Breeders are satisfied with the level of production and reproduction traits reached (Pavlík *et al.*, 2017). With respect to appropriate effective population size, it is important that more than 300 Slovak Dairy sires are produced each year. In the future, not only the numbers of Slovak Dairy population, its production and reproduction traits, but also its exterior traits, including wool characteristics, need to be continuously stabilized (Margetín, 2017c).

ACKNOWLEDGEMENTS

The study was supported by the projects APVV-0458-10, VEGA 1/0364/15 and KEGA 035 SPU-4/2015.

REFERENCES

- APOLEN, D. – ČAPISTRÁK, A. – MARGETÍN, M. 2000. Mlieková úžitkovosť a plodnosť kríženiak plemena merino s dojnými plemenami. *Acta Fytotechnica et Zootechnica*. Nitra: XIX. Dni genetiky, vol. 3, 2000, p. 121.
- ČAPISTRÁK, A. – MARGETÍN, M. – APOLEN, D. – ŠPÁNIK, J. 2000. Produkcia mlieka a plodnosť oviec plemena zošľachtená valaška, kríženiak s plemenami lacaune a východofrízskym plemenom. *Acta Fytotechnica et Zootechnica*. Nitra: XIX. Dni genetiky, vol. 3, 2000, p. 120.
- ČAPISTRÁK, A. – MARGETÍN, M. – APOLEN, D. – ŠPÁNIK, J. 2002. Produkcia a obsah základných zložiek mlieka oviec plemena zošľachtená valaška, lacaune a ich kríženiak. *Journal of Farm Animal Science*, vol. 35, 2002, p. 89–96.
- ČAPISTRÁK, A. – MARGETÍN, M. – APOLEN, D. – ŠPÁNIK, J. 2005. Produkcia a zloženie mlieka bahnic rôznych plemien a ich kríženiak počas dojnej periódy. *Journal of Farm Animal Science*, vol. 38, 2005, p. 181–189.
- ČAPISTRÁK, A. – MARGETÍN, M. – APOLEN, D. – ŠPÁNIK, J. 2006a. Porovnanie morfológických ukazovateľov vemena rôznych genotypov oviec. *Acta Fytotechnica et Zootechnica*, vol. 9, mimoriadne vydanie, 2006a, p. 179–181.
- ČAPISTRÁK, A. – MARGETÍN, M. – ŠPÁNIK, J. – APOLEN, D. 2006b. Lineárne hodnotenie, miery vemena a produkcia mlieka rôznych genotypov oviec. In: *Zborník prednášok „Ovce – Kozy Seč 2006“*, 10.–11.11.2006, p. 106–113. ISSN 1213–600X.
- MAČUHOVÁ, L. – UHRINČAĎ, M. – MAČUHOVÁ, J. – MARGETÍN, M. – TANČIN, V. 2008. The first observation of milkability of the sheep breeds Tsigai, Improved Valachian and their crosses with Lacaune. *Czech Journal of Animal Science*, vol. 53 (12), 2008, p. 528–536.
- MAČUHOVÁ, L. – UHRINČAĎ, M. – MAČUHOVÁ, J. – TANČIN, V. 2009. Milkability of Tsigai, Improved Valachian, and their crosses with Lacaune. *Acta Fytotechnica et Zootechnica*, vol. 12, spec. iss., 2009, p. 385–394.
- MAKOVICKÝ, P. – NAGY, M. – MAKOVICKÝ, P. 2013. Comparison of external udder measurements of the sheep breeds Improved Valachian, Tsigai, Lacaune and their crosses. *Chilean Journal of Agricultural Research*, vol. 73 (4), 2013, p. 366–371.
- MAKOVICKÝ, P. – NAGY, M. – MAKOVICKÝ, P. 2014. The comparison of ewe udder morphology traits of Improved Valachian, Tsigai, Lacaune breeds and their crosses. *Mljekarstvo*, vol. 64 (2), 2014, p. 86–93.
- MAKOVICKÝ, P. – MARGETÍN, M. – MAKOVICKÝ, P. 2015a. Genetic parameters for the linear udder traits of nine dairy ewes – short communication. *Veterinarski Archiv*, vol. 85 (5), 2015a, p. 577–582.
- MAKOVICKÝ, P. – MARGETÍN, M. – MILERSKI, M. 2015b. Estimation of udder cistern size in dairy ewes by ultrasonography. *Mljekarstvo*, vol. 65 (3), 2015b, p. 210–218.
- MAKOVICKÝ, P. – MILERSKI, M. – MARGETÍN, M. – MAKOVICKÝ, P. – NAGY, M. 2015c. Genetic parameters for the size of udder cisterns in ewes diagnosed by ultrasonography among breeds: Improved Valachian, Tsigai, Lacaune and their crosses. *Archivos de Zootecnia*, vol. 64 (248), 2015c, p. 403–408.
- MAKOVICKÝ, P. – MARGETÍN, M. – MAKOVICKÝ, P. 2017a. Estimation of genetic and phenotypic parameters for udder morphology traits in different dairy sheep genotypes. *Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis*, vol. 65 (1), 2017a, p. 105–110.
- MAKOVICKÝ, P. – MARGETÍN, M. – NAGY, M. – MAKOVICKÝ, P. 2017b. Growth curves in lambs of various genotypes created on the basis of Improved Valachian and Tsigai breeds. *Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis*, vol. 65 (1), 2017b, p. 111–118.
- MARGETÍN, M. 2016. Aktuálny stav šľachtienia slovenskej dojnej ovce. *Chov oviec a kôz*, roč. 36 (4), 2016, p. 27–37.
- MARGETÍN, M. 2017a. Nové národné plemeno oviec – slovenská dojná ovca. *Pol'nohospodár*, roč. 61 (10), 2017a, p. 2.
- MARGETÍN, M. 2017b. Slovenská dojná ovca – praktický výsledok inovačného procesu. *Agromagazín*, roč. 19 (7), 2017b, p. 36–37.
- MARGETÍN, M. 2017c. Slovenská dojná ovca – ako ďalej v šľachtení. *Chov oviec a kôz*, roč. 37 (2), 2017, p. 3–4.
- MARGETÍN, M. – ČAPISTRÁK, A. – KALIŠ, M. 1993. Produkcia mlieka cigájskych bahnic a kríženiak s východofrízskym plemenom vo vzťahu k hmotnosti jahniat v odchovej fáze. *Journal of Farm Animal Science*, vol. 26, 1993, p. 181–188.
- MARGETÍN, M. – ČAPISTRÁK, A. 1994. Zámery v šľachtiteľskom programe chovu oviec, kontrole úžitkovosti a dedičnosti. In: *Predpoklady efektívneho chovu oviec v súčasnosti*. Trenčín: Výskumný ústav živočíšnej výroby, Stanica chovu a šľachtienia oviec a kôz, 1994, p. 8–19.
- MARGETÍN, M. – APOLEN, D. – ČAPISTRÁK, A. 1999a. Zošľacht'ovanie merinských oviec v smere mäsovo – mliekovom. *Náš chov*, vol. 59 (5), 1999a, p. 17–19.

- MARGETÍN, M. – ČAPISTRÁK, A. – ŠPÁNIK, J. – APOLEN, D. – HYŽOVÁ, A. 1999b. Produkcia, zloženie mlieka a laktáčnité krivky bahnic plemena zošľachtená valaška, lacaune a ich kríženík v období cicania jahniat. *Journal of Farm Animal Science*, vol. 32, 1999b, p. 111–119.
- MARGETÍN, M. – ČAPISTRÁK, A. – APOLEN, D. 2000a. Program zošľachtovania merinských oviec s použitím plemena lacaune a východofrízskoho plemena v šľachtiteľsko-experimentálnych chovoch. *Chov oviec a kôz*, roč. 21 (1-2), 2000a, p. 24–27.
- MARGETÍN, M. – ČAPISTRÁK, A. – APOLEN, D. 2000b. Program zošľachtovania cigájskych oviec s použitím plemena lacaune a východofrízskoho plemena v šľachtiteľsko-experimentálnych chovoch. *Chov oviec a kôz*, vol. 21 (1-2), 2000b, p. 27–30.
- MARGETÍN, M. – ČAPISTRÁK, A. – APOLEN, D. 2000c. Program zošľachtovania valašských oviec s použitím plemena lacaune a východofrízskoho plemena v šľachtiteľsko-experimentálnych chovoch. *Chov oviec a kôz*, vol. 21 (3), 2000c, p. 4–7.
- MARGETÍN, M. – BULLOVÁ, M. – ČAPISTRÁK, A. 2004a. Rastové krivky jahniat rôznych genotypov vytvorených na báze plemena zošľachtená valaška a cigája. In: *Aktuálne problémy riešené v agrokomplexe* [CD]. Nitra: Slovenská poľnohospodárska univerzita v Nitre, 2004a, p. 456–461. ISBN 80-8069-488-6.
- MARGETÍN, M. – ČAPISTRÁK, A. – ŠPÁNIK, J. – APOLEN, D. – BULLOVÁ, M. 2004b. Rastová intenzita jahniat pri tvorbe rôznych genotypov plemena cigája. In: *Chov hospodárskych zvierat v podmienkach EÚ* [CD]. Nitra: Slovenská poľnohospodárska univerzita v Nitre, 2004b, p. 331–337.
- MARGETÍN, M. – ČAPISTRÁK, A. – APOLEN, D. – ORAVCOVÁ, M. – ŠPÁNIK, J. 2005a. Doterajšie výsledky programu zošľachtovania dojnych oviec na Slovensku. In: *Zborník z medzinárodnej vedeckej konferencie „Ekologické a ekonomické aspekty využitia poľnohospodársky znevýhodnených plôch chovom malých preživavcov“*. Žilina: VÚM, 2005a, p. 115–122.
- MARGETÍN, M. – MILERSKI, M. – APOLEN, D. – ČAPISTRÁK, A. – ORAVCOVÁ, M. 2005b. Morphology of udder and milkability of ewes of Tsigai, Improved Valachian, Lacaune breeds and their crosses. In: *Conference on „Physiological and technical aspects of machine milking*. Nitra: Slovak Republic, 26. – 28. apríl 2005b, ICAR Technical series no. 10, 2005b, p. 259–263. ISSN 1563-2504, ISBN 92-95014-07-3.
- MARGETÍN, M. – ŠPÁNIK, J. – MILERSKI, M. – ČAPISTRÁK, A. – APOLEN, D. 2005c. Relationships between morphological and functional udder traits and somatic cell count in milk of ewes. In: *Conference on „Physiological and technical aspects of machine milking*. Nitra: Slovak Republic, 26. – 28. apríl 2005, ICAR Technical series no. 10, 2005c, p. 255–258. ISSN 1563-2504, ISBN 92-95014-07-3.
- MARGETÍN, M. – MAKOVICKÝ, P. – ČAPISTRÁK, A. – APOLEN, D. – ORAVCOVÁ, M. – TANČIN, V. 2008. Odhad genetických parametrov vybraných ukazovateľov charakterizujúcich morfológiu vemena a dojitelnosť bahnic. In: *CD – zborník z medzinárodnej vedeckej konferencie „Aktuálne problémy riešené v Agrokomplexe“*, 5. 12. 2008, Nitra: SPU, 2008, p. 234–240, ISBN 978-80-552-0151-1.
- MARGETÍN, M. – APOLEN, D. – ČAPISTRÁK, A. – ŠPÁNIK, J. – MARGETÍNOVÁ, J. 2010. Postup šľachtenia syntetickej populácie oviec vytvorenej s použitím plemena lacaune a východofrízske. In: *Rozvoj metód zlepšovania a efektívnejšieho využívania genofondu zvierat z hľadiska konkurencieschopnosti, kvality produktov a ich zdravotnej bezpečnosti*. Nitra: Centrum výskumu živočíšnej výroby Nitra – Ústav chovu a šľachtovania oviec a kôz Trenčianska Teplá, 2010, p. 57.
- MARGETÍN, M. – APOLEN, D. – ČAPISTRÁK, A. – KULINOVÁ, K. 2011a. Results of improvement program in dairy sheep raised in Slovakia with using of Lacaune and East Friesian breed. In: *Možliwosci poprawy efektywnosci chowu owiec i kôz w Polsce w ujęciu regionalnym*. Kraków: Instytut Zootechniki, 2011a, p. 59. ISBN 978-83-7607-188-6.
- MARGETÍN, M. – APOLEN, D. – ČAPISTRÁK, A. – MILERSKI, M. – ORAVCOVÁ, M. – DEBRECÉNI, O. 2011b. Vplyv špecializovaných dojnych plemien na veľkosť cisterien vemena kríženík valašských oviec. *Acta Fytotechnica et Zootechnica*, vol. 13 (4), 2011b, p. 85–92.
- MARGETÍN, M. – ČAPISTRÁK, A. – APOLEN, D. – RELOVSKÝ, S. – ŠUTÝ, J. – RAFAJOVÁ, M. 2011c. Tvorba syntetickej populácie slovenskej dojnej ovce – druhá etapa programu zošľachtovania. *Chov oviec a kôz*, vol. 21 (4), 2011c, p. 6–8.
- MARGETÍN, M. – MAKOVICKÝ, P. – MILERSKI, M. – APOLEN, D. – DEBRECÉNI, O. – ORAVCOVÁ, M. 2011d. The effect of specialized dairy breeds on udder cistern size in Tsigai crossbreeds. *Slovak Journal of Animal Science*, vol. 44 (4), 2011, p. 146–153.
- MARGETÍN, M. – ORAVCOVÁ, M. – APOLEN, D. – MILERSKI, M. 2012a. Genetic parameters for udder traits in Slovak dairy sheep and their crosses with specialized breeds. *Journal of Life Sciences*, no. 6, 2012a, p. 1363–1370.
- MARGETÍN, M. – APOLEN, D. – ČAPISTRÁK, A. 2012b. Šľachtenie slovenskej dojnej ovce – druhá etapa zošľachtovacieho programu. *Slovenský chov*, vol. 17 (1), 2012b, p. 25–27.

- MARGETÍN, M. – MILERSKI, M. – APOLEN, D. – ČAPISTRÁK, A. – ORAVCOVÁ, M. – DEBRECÉNI, O. 2013a. Relationships between production, quality of milk and udder health status of ewes during machine milking. *Journal of Central European Agriculture*, vol. 14 (1), 2013a, p. 328–340.
- MARGETÍN, M. – ORAVCOVÁ, M. – MAKOVICKÝ, P. – APOLEN, D. – DEBRECÉNI, O. 2013b. Milkability of Improved Valachian, Tsigai and Lacaune purebred and crossbred ewes. *Slovak Journal of Animal Science*, vol. 46 (3), 2013b, p. 100–109.
- MARGETÍN, M. – BÍREŠ, J. – REĽOVSKÝ, S. – ORAVCOVÁ, M. – MOJŽIŠ, M. – RAJSKÁ, Z. 2016. Changes in allele and genotype frequencies of PrP gene in breeds of sheep in Slovakia between 2004 and 2015. *Acta Fytotechnica et Zootechnica*, vol. 19 (3), 2016, p. 119–122.
- MARGETÍN, M. – APOLEN, D. – ČAPISTRÁK, A. 2017. História procesu šľachtenia slovenskej dojnej ovce. *Slovenský chov*, vol. 22 (5), 2017, p. 14–17.
- MILERSKI, M. – MARGETÍN, M. – APOLEN, D. – ČAPISTRÁK, A. – ŠPÁNIK, J. 2005. Udder cistern size and milkability of ewes of various genotypes. In: *Conference on „Physiological and technical aspects of machine milking. Nitra: Slovak Republic, 26.–28. apríl 2005, ICAR Technical series no. 10, 2005. p. 63–69.*
- MILERSKI, M. – MARGETÍN, M. – ČAPISTRÁK, A. – APOLEN, D. – ŠPÁNIK, J. – ORAVCOVÁ, M. 2006. Relationships between external udder measurements and the linear scores for udder morphology traits in dairy sheep. *Czech Journal of Animal Science*, vol. 51 (9), 2006, p. 383–390.
- PAVLÍK, I. – MARGETÍN, M. – JANÍČEK, M. 2017. Slovenská dojná ovca – výsledky u popredných chovateľov sú prísľubom budúcnosti. *Chov oviec a kôz*, vol. 37 (2), 2017, p. 17–18.
- TANČIN, V. – MAČUHOVÁ, L. – ORAVCOVÁ, M. – UHRINČAŤ, M. – KULINOVÁ, K. – ROYCHOUDHURY, S. – MARNET, P. G. 2011. Milkability assessment of Tsigai, Improved Valachian, Lacaune and F1 Crossbred ewes (Tsigai x Lacaune, Improved Valachian x Lacaune) throughout lactation. *Small Ruminant Research*, vol. 97, 2011, p. 28–34.