

Short communication

THE EFFECT OF *CURCUMA LONGA* DRIED POWDER IN THE DIET ON WEIGHT GAIN OF RABBIT DOES

M. FÖLDEŠIOVÁ^{1*}, A. BALÁŽI², L. CHRASTINOVÁ², P. CHRENEK^{1,2}

¹Slovak University of Agriculture, Faculty of Biotechnology and Food Science, Nitra, Slovak Republic

²National Agricultural and Food Centre, Research Institute for Animal Production Nitra, Lužianky, Slovak Republic

ABSTRACT

The aim of the present study was to evaluate the effect of different concentrations of *Curcuma longa* plant additive to the diet on the weight gain of rabbit does. Rabbit females ($n = 45$) of New Zealand White breed of the same age (2 months) were used in the experiment. Rabbit does in the control group (C; $n = 15$) were fed a commercially available feed. In the experimental groups 5 g (E1; $n = 15$) and 20 g (E2; $n = 15$) of *Curcuma longa* dried powder was added to 100 kg feed mixture. The highest average weight gain per week was observed in the first experimental group (E1; 235.7 ± 22.35 g) when compared to the control (C; 216.2 ± 25.59 g) and the second experimental (E2; 220.5 ± 31.94 g) groups. The highest total average weight gain of rabbit does was observed in the E1 (2103.3 ± 63.22 g) compared to the second (E2, 2045 ± 84.36 g) and control (C, 1950 ± 126.88 g) groups. In conclusion, supplementation of *Curcuma longa* plant powder to the commercially diet for rabbit positively affected weight gains in rabbit does. Therefore, to improve growth performance, further studies are required to define an optimal supplementation of *Curcuma longa* to the rabbit diet.

Keywords: rabbit does; *Curcuma longa*; weight gain

INTRODUCTION

In the last years, rabbit production on the commercial level acquired increasing interest due to their prolificacy, rapid growth rate and meat yields (Savietto *et al.*, 2015; Mínguez, 2014; Ricke *et al.*, 2012; Gondret *et al.*, 2005).

More suitable composition of feed mixture or administration of natural additives at a suitable concentration might be beneficial in livestock farming without negative effect on the environment and the animal as an individual (Wareth *et al.*, 2014; Githiori *et al.*, 2003; Amber *et al.*, 2004; Namkung *et al.*, 2004).

Positive effect of biologically active substances and extracts from *Acacia saligna* (Tamir and Asefa, 2009), *Morus australis* (Wu *et al.*, 2013), *Yucca schidigera* (Földešiová *et al.*, 2013), *Agave tequilana* (Sáyago-Ayerdi *et al.*, 2014), *Saposhnikovia divaricata*, *Lonicera japonica*, *Chelidonium majus* (Park *et al.*, 2014) a. o. on weight gain were observed in lambs, mice, rabbits, rats and broilers.

Curcuma longa Linn, a member of *Zingiberaceae* family, commonly known as turmeric, originate in tropical and subtropical regions of India and China. Medicinal properties of *Curcuma longa* have been attributed primarily to curcuminoids, which are located

*Correspondence: e-mail: mt.foldesiova@gmail.com

Martina Földešiová, Institute of Farm Animal Genetics and Reproduction, Research Institute for Animal Production in Nitra, National Agricultural and Food Centre, Lužianky, Slovak Republic, phone: +421 37 6546285, Fax: +421 37 654 61 89

in the plant rhizome. Curcumin (diferuloylmethane) is the most important fraction of *Curcuma longa* (Araújo and Leon, 2001). Polyphenol curcumin, extracted from dried rhizomes of the plant, acts through inhibition of mitogen-activated protein kinases (Jeon *et al.*, 2013). Although turmeric is consumed every day in Asian countries, no toxic effect on the health of population was found (Ammon and Wahl, 1991).

Beneficial effects of curcuma on the animal liver protection (Deshpande *et al.*, 1998), treatment of the human diabetes (Eshrat and Hussain, 2002), steroidogenesis, proliferation and apoptosis in porcine ovaries (Kádasi *et al.*, 2012) were found.

In particular, yellow rhizome, containing curcumin, is used to treat digestive, neuropsychiatric diseases (Mei *et al.*, 2011) and osteoarthritis in combination with a ginger (Low Dog, 2006).

Furthermore, the plant has also been shown to possess high antioxidant (Maheshwari *et al.*, 2006; Wojdyło *et al.*, 2007) anticarcinogenic (Hatcher *et al.*, 2008), antibacterial (De *et al.*, 2009) and anti-inflammatory (Jurenka, 2009) effect.

The objective of this study was to evaluate the effect of different concentrations of *Curcuma longa* dried powder as an additive to the diet on the average weight gain (g) per week and the total average weight gain of the rabbit does.

MATERIAL AND METHODS

Animals

Two months old clinically healthy rabbit does of the New Zealand White line (NAFC Nitra, SR) were used in this experiment. The animals were housed in individual cages, under a constant photoperiod of 14 hours of light day, average relative humidity $60 \pm 5\%$ and temperature $17 \pm 3\text{ }^{\circ}\text{C}$. The rabbits were fed *ad libitum* and water was provided *ad libitum* with nipple drinkers.

Rabbit does ($n = 45$) were divided into three groups: control (C; $n = 15$) and two experimental groups (E1; $n = 15$ and E2; $n = 15$). The does in the control group were fed a commercially available complete feed mixture. In experimental groups the complete feed mixture was enriched with *Curcuma longa* dried powder at the concentrations of 5 g (E1) and 20 g (E2) per 100 kg. The animals were fed for 63 days (9 weeks) and weighted weekly.

The treatment of the animals was approved by the Ministry of Agriculture and Rural Development of the Slovak Republic, no. SK P 28004 and Ro 1488/06-221/3a.

Statistical analysis

The data were analysed by the t-test using Sigma Plot statistical package (Systat Software Inc., Germany).

RESULTS AND DISCUSSION

In our study we tested effect of the addition of *Curcuma longa* dried powder to the complete feed mixture on the average weight gain (g) per week and the total average weight gain (g) of rabbit does.

The highest average weight gain of rabbit does per week (g) was found in the first experimental group (E1; 235.7 ± 22.35) when compared to the control (C; 216.2 ± 25.59) and the second experimental group (E2; 220.5 ± 31.94) (Table 1).

Total average weight gain (g) was higher in the first experimental group (E1; 2103.3 ± 63.22) compared to the second (E2; 2045 ± 84.36) and control (C; 1950 ± 126.88) groups (Table 2, Figure 1).

In accordance to Holder *et al.* (1978) we suggest that slightly lower weight gain in the second experimental group compared to the first experimental group might be due to the higher concentration of *Curcuma longa* in feed mixture, which can cause poor absorption from the intestine.

In our study, we found that the addition of both concentrations (5 g and 20 g kg^{-1} diet) of *Curcuma longa* dried powder into rabbit complete feed mixture had a positive effect on average weight gain per week and total average weight gain of analysed rabbit does.

Positive effect of *Curcuma longa* powder to the diet was also found in broiler chickens. Higher weight gain was observed in the birds fed the diet containing *Curcuma longa* at level of 0.5 %, compared to the birds receiving 0.25 %, 1 % and control birds (Al-Sultan, 2003). Osava *et al.*, (1995) and Al-Sultan (2003) attributed the increase in the body weight gain to the antioxidant activity of *Curcuma longa*.

Moreover, Durrani *et al.* (2006) reported significantly positive effect of curcuma at the level of 0.5 % on weight gain of birds. It was also shown, that curcumin added to the diet of kids during the hot summer months significantly improved the final live body weight and average daily body gain of kids compared to the control (Habebband and Tarabany, 2012). On the other hand, similarly to our results no significant effect of the supplementation of curcuma powder in the broiler rabbit (Basavaraj *et al.*, 2010) and in the broiler chicks feed a mixture (Mehala and Moorthy, 2008) were reported.

Table 1: Weight gain (g) per week of analysed rabbit does fed with *Curcuma longa* enriched feed

Groups	Weight gain per week (g)										Average weight gain per week (g) (Mean \pm S.E.M.)
	1 st week	2 nd week	3 rd week	4 th week	5 th week	6 th week	7 th week	8 th week	9 th week	10 th week	
C ($n = 15$)	286.20 \pm 80.77	280.77 \pm 27.86	258.46 \pm 23.26	245.38 \pm 28.43	223.08 \pm 27.29	185.38 \pm 31.88	227.69 \pm 47.73	183.85 \pm 19.29	59.23 \pm 29.41	216.20 \pm 25.59	
E1 ($n = 15$)	258.00 \pm 29.13	291.30 \pm 17.75	310.70 \pm 19.28	283.30 \pm 21.86	278.00 \pm 29.18	217.30 \pm 28.07	208.00 \pm 22.28	170.70 \pm 20.31	86.00 \pm 17.91	235.70 \pm 22.35	
E2 ($n = 15$)	222.86 \pm 27.08	263.57 \pm 30.69	327.10 \pm 40.35	299.30 \pm 16.82	254.30 \pm 18.53	232.90 \pm 14.76	211.40 \pm 22.84	165.00 \pm 32.39	69.30 \pm 27.60	220.50 \pm 31.94	

C – control group, commercially available diet (normal diet); E1 – 5 g of *Curcuma longa* dried powder added to 100 kg of normal diet; E2 – 20 g of *Curcuma longa* dried powder added to 100 kg of normal diet

Table 2: Average weight per week (g) and total average weight gain of analysed rabbit does fed with *Curcuma longa* enriched feed

Groups	Average weight of rabbit does per week (g)										Total average weight gain (g) (Mean \pm S.E.M.)
	1 st week	2 nd week	3 rd week	4 th week	5 th week	6 th week	7 th week	8 th week	9 th week	10 th week	
C ($n = 15$)	1713.08 \pm 83.30	1999.23 \pm 103.41	2280.00 \pm 116.59	2538.46 \pm 128.04	2783.85 \pm 148.68	3006.92 \pm 150.51	3192.31 \pm 156.99	3420.00 \pm 183.50	3603.85 \pm 187.85	3663.08 \pm 171.73	1950.00 \pm 126.88
E1 ($n = 15$)	1842.67 \pm 44.37	2100.67 \pm 53.24	2392.00 \pm 58.98	2702.67 \pm 65.03	2986.00 \pm 64.66	3264.00 \pm 72.93	3481.33 \pm 73.03	3689.33 \pm 86.68	3860.00 \pm 78.22	3946.00 \pm 77.07	2103.30 \pm 63.22
E2 ($n = 15$)	1818.57 \pm 47.31	2041.43 \pm 57.87	2305.00 \pm 63.48	2632.14 \pm 65.84	2931.43 \pm 77.90	3185.71 \pm 77.78	3418.57 \pm 86.23	3630.00 \pm 102.61	3795.00 \pm 118.74	3864.29 \pm 104.18	2045.00 \pm 84.36

C – control group, commercially available diet (normal diet); E1 – 5 g of *Curcuma longa* dried powder added to 100 kg of normal diet; E2 – 20 g of *Curcuma longa* dried powder added to 100 kg of normal diet




Fig. 1: Total average weight gain (g) of analysed rabbit does fed with *Curcuma longa* dried plant added into complete feed mixture

CONCLUSION

The supplementation of *Curcuma longa* plant powder to the commercially available diet for rabbits positively affects weight gains in rabbit does. Therefore, for improving growth performance, further studies are required to define an optimal supplementation of *Curcuma longa* to the rabbit diet.

ACKNOWLEDGMENT

We thank Mr. Milan Dobias̄ for technical support. The research leading to these results has received funding from the European Community under project no 26220220180: Building Research Centre „AgroBioTech“.

REFERENCES

- AL-SULTAN, S. I. 2003. The Effect of *Curcuma longa* (Turmeric) on Overall Performance of Broiler Chickens. *International Journal of Poultry Science*, vol. 2 (5), 2003, p. 351–353.
- AMBER, K. H. – YAKOUT, H. M. – HAMED RAWYA, S. 2004. Effect of feeding diets containing yucca extract or probiotic on growth, digestibility, nitrogen balance and caecal microbial activity of growing New Zealand White rabbits. In: *Proceedings – 8th*

World Rabbit Congress. Puebla, Mexico, p. 737–745.

- AMMON, H. P. – WAHL, M. A. 1991. Pharmacology of *Curcuma longa* L. *Planta Medica*, vol. 57, 1991, p. 1–7.
- ARAÚJO, C. A. C. – LEON, L. L. 2001. Biological activities of *Curcuma Longa* L. *Memorias Do Instituto Oswaldo Cruz*, vol. 96 (5), 2001, p. 723–728.
- BASAVARAJ, M. – NAGABHUSHANA, V. – PRAKASH, N. – MALLIKARJUNAPPA, S. – APPANNAVAR, M. M. – WAGMARE, P. 2010. Effect of dietary supplementation of Pulvis *Curcuma Longa* on the voluntary feed intake, nutrient digestibility and growth performance of Broiler rabbits under summer stress. *Veterinary World*, vol. 3 (8), 2010, p. 369–372.
- DE, R. – KUNDU, P. – SWARNAKAR, S. – RAMAMURTHY, T. – CHOWDHURY, A. – NAIR, G.B. – MUKHOPADHYAY, A. K. 2009. Antimicrobial Activity of Curcumin against *Helicobacter pylori* Isolates from India and during Infections in Mice. *Antimicrobial Agents and Chemotherapy*, vol. 53 (4), 2009, p. 1592–1597.
- DESHPANDE, U.R. – GADRE, S. G. – RASTE, A. S. – PILLAI, D. – BHIDE, S. V. – SAMUEL, A. M. 1998. Protective effect of turmeric (*Curcuma longa* L.) extract on carbon tetrachloride-induced liver damage in rats. *Indian Journal of Experimental Biology*, vol. 36 (6), 1998, p. 573–577.
- DURRANI, F.R. – ISMIAL, M. – SULTAN, A. – SHHAIL, S. M. – CHAND, N. – DURRANI, Z. 2006. Effect of different levels of feed added turmeric

- (*Curcuma longa*) on the performance of broiler chicks. *Journal of Agricultural and Biological Science*, vol. 1 (2), 2006, p. 9–11.
- ESHMAT, H. – HUSSAIN, A. 2002. Hypoglycemic, hypolipidemic and antioxidant properties of combination of Curcumin from *Curcuma longa*, Linn, and partially purified product from *Abroma augusta*, Linn. In streptozotocin induced diabetes. *Indian Journal of Clinical Biochemistry*, vol. 17 (2), 2002, p. 34–43.
- FÖLDEŠIOVÁ, M. – BALÁŽI, A. – CHRASTINOVÁ, E. – CHRENEK, P. 2013. Effect of *Yucca schidigera* herbal extract in diet on rabbit does weight gains. *Slovak Journal of Animal Science*, vol. 46 (2), 2013, p. 81–86.
- GITHIORI, J. B. – HÖGLUND, J. – WALLER, P. J. – LEYDEN BAKER, R. 2003. Evaluation of antihelmintic properties of extracts from some plants used as livestock dewormers by pastoralist and smallholder farmers in Kenya against *Heligmosomoides polygyrus* infections in mice. *Veterinary Parasitology*, vol. 118 (3–4), p. 215–226.
- GONDRET, F. – LARZUL, C. – COMBES, S. – DE ROCHAMBEAU, H. 2005. Carcass composition, bone mechanical properties, and meat quality traits in relation to growth rate in rabbits. *Journal of Animal Science*, vol. 83 (7), p. 1526–1535.
- HABEEBAND, A. A. M. – EL-TARABANY, A. A. 2012. Effect of *Nigella sativa* or Curcumin on daily body weight gain, feed intake and some physiological functions in growing Zaraibi Goats during hot summer season. *Arab Journal of Nuclear Science and Applications*, vol. 45 (2), 2012, p. 238–249.
- HATCHER, H. – PLANALP, R. – CHO, J. – TORTI, F. M. – TORTI, S. V. 2008. Curcumin: From ancient medicine to current clinical trials. *Cellular and Molecular Life Sciences*, vol. 65 (11), 2008, p. 1631–1652.
- HOLDER, G.M. – PLUMMER, J.L. – RYAN, A. J. 1978. The metabolism and excretion of curcumin (1,7-bis-(4-hydroxy-3-methoxyphenyl)-1,6-hepadiene-3,5-dione) in the rat. *Xenobiotica*, vol. 8 (12), 1978, p. 761–768.
- JEON, Y. – KIM, CH. E. – JUNG, D. – KWAK, K. – PARK, S. – LIM, D. – KIM, S. – BAEK, W. 2013. Curcumin Could Prevent the Development of Chronic Neuropathic Pain in Rats with Peripheral Nerve Injury. *Current Therapeutic Research*, vol. 74, 2013, p. 1–4.
- JURENKA, J.S. 2009. Anti-inflammatory Properties of Curcumin, a Major Constituent of *Curcuma longa*: A Review of Preclinical and Clinical Research. *Alternative Medicine Review*, vol. 14 (2), 2009, p. 141–153.
- KÁDASI, A. – SIROTKIN, A. V. – MARUNIAKOVÁ, N. – KOLESÁROVÁ, A. – BULLA, J. – GROSSMAN, R. 2012. The effect of curcumin on secretory activity, proliferation and apoptosis of the porcine ovarian granulosa cells. *Journal of Microbiology, Biotechnology and Food Sciences*, vol. 2 (1), 2012, p. 349–357.
- LOW DOG, T. 2006. A Reason to Season: The Therapeutic Benefits of Spices and Culinary Herbs. *The Journal of Science and Healing*, vol. 2 (5), 2006, p. 446–449.
- MAHESHWARI, R. K. – SINGH, A. K. – GADDIPATI, J. – SRIMAL, R. C. 2006. Multiple biological activities of curcumin: A short review. *Life Sciences*, vol. 78 (18), 2006, p. 2081–2087.
- MEI, X. – XU, D. – XU, S. – ZHENG, Y. – XU, S. 2011. Gastroprotective and antidepressant effects of new zinc (II)-curcumin complex in rodent models of gastric ulcer and depression induced by stresses. *Pharmacology, Biochemistry and Behavior*, vol. 99 (1), 2011, p. 66–74.
- MEHALA, C. – MOORTHY, M. 2008. Production performance of broilers fed with Aloe vera and *Curcuma longa* (Turmeric). *International Journal of Poultry Science*, vol. 7 (9), 2008, p. 852–856.
- MÍNGUEZ, B. C. 2014. Genetic analyses of growth, carcass and meat quality traits in maternal lines of rabbits and their diallel cross: tesis doctoral. Valencia : Universitat Politècnica de Valencia. 261s.
- NAMKUNG, H. – LI, M. – GONG, J. – YU, H. – COTTRILL, M. – DE LANGE, C. F. M. 2004. Impact of feeding blends of organic acids and herbal extracts on growth performance, gut microbiota and digestive function in newly weaned pigs. *Canadian Journal of Animal Science*, vol. 84 (4), p. 697–704.
- OSAWA, T. – SUGIYAMA, Y. – INAYOSHI, M. – KAWAKISHI, S. 1995. Antioxidative activity of tetrahydrocurcuminoids. *Bioscience, Biotechnology and Biochemistry*, vol. 59 (9), p. 1609–1612.
- PARK, J. H. – KANG, S. N. – CHU, G. M. – JIN, S. K. 2014. Growth performance, blood cell profiles, and meat quality properties of broilers fed with *Saposhnikovia divaricata*, *Lonicera japonica* and *Chelidonium majus* extracts. *Livestock Science*, vol. 165, 2014, p. 87–94.
- RICKE, S. C. – VAN LOO, E. J. – JOHNSON, M. G. – O'BRYAN, C. A. 2012. *Organic Meat Production and Processing*. The first edition. USA : Wiley-Blackwell. 464 p. ISBN 978-0-8138-2126-9.
- SAVIETTO, D. – FRIGGENS, N. C. – PASCUAL, J. J. 2015. Reproductive robustness differs between generalist and specialist maternal rabbit lines: the role of acquisition and allocation of resources. *Genetics, Selection, Evolution*, vol. 47, 2015, p. 1–12.

- SÁYAGO-AYERDI, S. – MATEOS, R. – ORTIZ-BASURTO, R. I. – LARGO, C. – SERRANO, J. – BELÉN GRANADO-SERRANO, A. – SARRIÁ, B. – BRAVO, L. – TABERNERO, M. 2014. Effects of consuming diets containing *Agave tequilana* dietary fibre and jamaica calyces on body weight gain and redox status in hypercholesterolemic rats. *Food Chemistry*, vol. 148 (1), 2014, p. 54–59.
- TAMIR, B. – ASEFA, G. 2009. Effects of different forms of *Acacia saligna* leaves inclusion on feed intake, digestibility and body weight gain in lambs fed grass hay basal diet. *Animal Feed Science and Technology*, vol. 153 (1–2), 2009, p. 39–47.
- ABDEL-WARETH, A. A. A. – HAMMAD, S. – AHMED, H. 2014. Effects of *Khaya Senegalensis* leaves on performance, carcass traits, haematological and biochemical parameters in rabbits. *EXCLI Journal*, vol. 13, 2014, p. 502–512.
- WOJDYŁO, A. – OSZMIANSKI, J. – CZEMERYŚ, R. 2007. Antioxidant activity and phenolic compounds in 32 selected herbs. *Food Chemistry*, vol. 105 (3), 2007, p. 940–949.
- WU, T. – QI, X. – LIU, Y. – GUO, J. – ZHU, R. – CHEN, W. – ZHENG, X. – YU, T. 2013. Dietary supplementation with purified mulberry (*Morus australis* Poir) anthocyanins suppresses body weight gain in high-fat diet fed C57BL/6 mice. *Food Chemistry*, vol. 141 (1), 2013, p. 482–487.