



Investigations on the influence of roughage/concentrate ratio and linseed oil supplementation on rumen fermentation and microbial protein yield in dairy cows

G. FLACHOWSKY, K. ERDMANN, P. LEBZIEN, L. HÜTHER

Institute of Animal Nutrition of the Federal Agricultural Research Centre, Germany

ABSTRACT

The objective of the study was to investigate the influence of the roughage/concentrate ratio and linseed oil supplementation on rumen fermentation and microbial protein yield in late lactating dairy cows.

Seven late lactating double fistulated cows were randomly assigned to four experimental periods applying an incomplete 2 x 2 Latin square design. The rations consisted of meadow hay and a concentrate mixture given in a ratio of 70 : 30 (H 70) or 30 : 70 (H 30) on a dry matter basis. The basal mixtures fed were either supplemented with 200 g linseed oil daily (HLO 70 and HLO 30) or non-supplemented. After about three weeks of adaptation to the experimental rations, samples from the rumen and the duodenal chyme were taken to study parameters of rumen fermentation and the nutrient flow to the duodenum.

The pH-value, the NH₃-N concentration, and the content of short-chain fatty acids in the rumen as well as the apparent ruminal digestibility of organic matter (OM) and neutral detergent fibre (NDF) were not significantly affected by linseed oil supplements. However, low portions of hay in the ration significantly reduced the pH-value and increased the NH₃-N concentration.

An increased hay portion in the ration resulted in an increased NDF-degradation in the rumen as compared to the rations high in concentrate. Due to an increased flow of microbial OM after high concentrate feeding, the "apparent ruminal digestibility" of OM was calculated to decrease although the quantity of FOM (fermented organic matter) increased. Feeding the rations poor in hay increased the flow of non ammonia nitrogen (NAN) and utilisable crude protein (uCP) significantly due to the significantly increased microbial protein synthesis (g MP/d) and a improved efficiency of microbial protein synthesis (g MP/kg FOM) as well.

Keywords: dairy cows, roughage: concentrate, linseed oil, rumen fermentation, microbial protein yield

INTRODUCTION

The health and performance of dairy cows depends on the intake of digestible and metabolisable nutrients. Certain amounts of fibre from roughage are necessary for rumination and rumen processes (e.g. de Brabander et al., 1999; GfE, 2001; Meyer et al., 2001; 2002; Piatkowski, 1977). But normally roughage is less digestible than concentrate and can not provide enough energy for high yielding cows.

Fat can be added as so-called ruminally-protected or not protected fat. Unprotected fat supplements may affect rumen fermentation (Machmüller and Kreuzer, 1999; Palmquist and Kenkins, 1980; Sutton et al., 1983) and microbial protein synthesis (Doreau and Ferlay,

1995). Furthermore, added fats can influence fatty acids pattern of milk fat (e.g. Flachowsky et al., 1996; Griinari and Bauman, 2006; Jahreis et al., 1996; Kennely and Glimm, 1998; Wagner et al., 1998) depending on the fatty acid pattern of the supplemented fats.

Fewer studies are available investigating the effects of oils with high concentrations of unsaturated fatty acids, such as linseed oil, on parameters of rumen fermentation, and nutrient flow to the duodenum of cows fed with various kinds of diets. Therefore, the objective of the present study was to investigate the influence of two roughage/concentrate ratios (70 : 30 and 30 : 70) with and without linseed oil supplementation on parameters of rumen fermentation and nutrient flow to the duodenum in late lactating dairy cows.

This article is dedicated to our friend Prof. Dr. Alexander Sommer on the occasion of his 70th birthday

Correspondence: E-mail: gerhard.flachowsky@fal.de

MATERIAL AND METHODS

Seven multiparous lactating Friesian cows (initial body weight: 554 ± 45 kg) fitted with cannulae in the dorsal rumen, and simple T-shaped cannulae in the proximal duodenum, were randomly assigned to four experimental periods applying an incomplete 2×2 Latin square design. The cows were in late lactation (230 ± 25 days in milk), producing 21.8 ± 4.1 kg fat corrected milk per day. The animals were kept individually on rubber mats with slatted flooring without bedding material. Drinking water was freely available.

The cows were fed with four different diets varying in percentages of meadow hay (70 and 30 % of DM) and concentrate (30 and 70 % of DM) both supplemented with 200 g linseed oil per cow per day or unsupplemented (Table 1).

Table 1: Feeding of cows in various periods

| Feedstuffs | Ration | | | |
|---|--------|--------|------|--------|
| | H 70 | HLO 70 | H 30 | HLO 30 |
| Meadow hay ¹⁾ (% of DMI) | 70 | 70 | 30 | 30 |
| (kg DM day ⁻¹) | 9.1 | 9.1 | 4.0 | 4.0 |
| Concentrate ²⁾ (% of DMI) | 30 | 30 | 70 | 70 |
| (kg DM day ⁻¹) | 3.9 | 3.9 | 8.9 | 8.9 |
| Linseed oil (g d ⁻¹) | - | 200 | - | 200 |

¹⁾ 98.5 g crude protein, 625 g NDF per kg DM,

²⁾ Ingredients of concentrate: 38% barley, 35% wheat, 13% rye, 11.5% extr. soybeanmeal, 2.5% vit/min premix; 177.0 g crude protein, 214.0 g NDF per kg DM

To avoid feed refusals, all cows consumed adequate amounts of feed based on the dry matter intake (DMI) of the cow with the lowest DMI. Concentrate was given in four (equally-sized) portions at 5:30 and 7:30 a.m. and 4:30 and 6:30 p.m. Meadow hay was offered at 5:30 a.m. and 4:30 p.m. At 5:30 a.m. and 4:30 p.m., 100 g of linseed oil (Protein- und Ölwerk Neuss GmbH & Co KG, Neuss, Germany) were administered directly onto the feedstuffs with a syringe.

Diets were given to the cows for four weeks, three weeks were planned for adaptation, chymus collection was carried out in the fourth week. Cows were milked at 5:00 a.m. and 3:00 p.m. and milk yield was recorded at each milking.

Ruminal fluid for the determination of pH, ammonia and short chain fatty acids (SCFA) was sampled during the third week for two days via the rumen cannulae using a hand vacuum pump at 5:30, 6:00, 6:30, 7:00, 7:30, 8:30 and 10:30 a.m. Rumen pH was measured immediately after sampling using a digital pH meter (pH 525, WTW, Weilheim, Germany). The ammonia content of rumen fluid was determined according to DIN 38406-E5-2 (1998).

The concentrations of SCFA were analysed by gas chromatography (Hewlett Packard 5580; Hewlett Packard, Avondale, PA, USA) with a flame ionization detector after sample-preparation according to Geisler et al. (1976). Duodenal flow was measured during the last week of each experimental period. To estimate the digesta flow, Cr_2O_3 was mixed with wheat flour, weighed into filter paper and given in two portions per day into the rumen beginning ten days before the start of the duodenal digesta collection, and in four portions per day during the collection period (Rohr et al., 1984). Spot samples of duodenal chyme were taken in 2-h intervals from Monday to Saturday as described by Rohr et al. (1979). After nitrogen (N; Kjeldahl method) determination in the fresh duodenal digesta, the digesta was freeze-dried and ground to pass through a 1mm-sieve. Cr_2O_3 in marker and digesta samples was analysed applying atomic absorption spectrometry according to Williams et al. (1962). The microbial portion of non-ammonia-N (NAN) in the freeze-dried duodenal samples was estimated using near infra-red spectroscopy (NIRS) according to Lebzién and Paul (1997). The chemical composition of the feedstuffs, as well as the NDF content of the digesta samples were determined according to the VDLUFA methods (Naumann and Bassler, 1993).

RESULTS AND DISCUSSION

The dry matter intake (DMI) of cows fed with diets high and low in roughage or concentrate and supplemented with 200 g linseed oil per day or non-supplemented was nearly the same (Table 2).

Table 2: Dry matter, nutrients and energy intake of cows during experimental periods

| Parameters | Ration | | | |
|------------------|------------------------------|-------------------------------|-------------------------------|------------------------------|
| | H 70 | HLO 70 | H 30 | HLO 30 |
| | kg day ⁻¹ | | | |
| DM | 12.82 ± 0.08 | 12.94 ± 0.35 | 12.83 ± 0.03 | 13.04 ± 0.18 |
| OM | 12.05 ^a ± 0.03 | 12.17 ^{ab} ± 0.28 | 12.19 ^{ab} ± 0.03 | 12.38 ^b ± 0.07 |
| CP | 1.52 ^a ± 0.06 | 1.61 ^a ± 0.12 | 2.01 ^b ± 0.13 | 1.98 ^b ± 0.13 |
| Ether extract | 0.26 ^a ± 0.02 | 0.42 ^b ± 0.04 | 0.29 ^a ± 0.10 | 0.47 ^b ± 0.04 |
| NDF | 6.45 ^b ± 0.07 | 6.39 ^b ± 0.19 | 4.36 ^a ± 0.08 | 4.39 ^a ± 0.10 |
| | MJ day ⁻¹ | | | |
| ME | 126.5 ^a ± 0.4 | 131.3 ^b ± 3.3 | 148.5 ^c ± 0.4 | 154.4 ^d ± 0.8 |
| NEL | 76.3 ^a ± 0.3 | 79.3 ^b ± 2.1 | 93.8 ^c ± 0.2 | 97.6 ^d ± 0.4 |

^{a, b, c, d} = values in the rows with different letters differ significantly ($p \leq 0.05$)

Cows fed with low hay (H 30 and HLO 30) consumed less fibre, but more protein and energy. Oil supplementation (HLO 70 and HLO 30) increased the intake of ether extract and energy in comparison to the non-supplemented groups (Table 2).

Rumen fermentation

The pH-value in the rumen liquid was significantly influenced by ration composition, but mostly not influenced ($p > 0.05$) by the supplementation of linseed oil (Table 3).

value (Table 3). The data are in agreement with references, where no significant effects of linseed oil supplementation on rumen pH were described (e.g. Doreau and Ferlay, 1995; Ueda et al., 2003). Doreau et al. (1993) added 0.99 kg rapeseed oil per day to rations of dairy cows and did not observe any influence on rumen pH-value.

The ammonia-concentration in the rumen increased up to 90 – 120 min after feeding (Table 4) and later decreased.

Both 30 and 60 minutes after feeding, the $\text{NH}_3\text{-N}$ -concentration was significantly higher in animals fed

Table 3: Effects of feeding on the pH-value in rumen liquid before and after morning feeding

| Time after feeding (min) | Ration | | | | | Effects ¹⁾ | | |
|--------------------------|--------|--------|------|--------|------|-----------------------|-------------|-------|
| | H 70 | HLO 70 | H 30 | HLO 30 | PSEM | Hay | Linseed oil | HxLO |
| Before | 6.55 | 6.49 | 6.88 | 6.81 | 0.07 | <0.001 | 0.424 | 0.938 |
| 30 | 6.47 | 6.37 | 6.69 | 6.57 | 0.08 | 0.014 | 0.177 | 0.902 |
| 60 | 6.37 | 6.22 | 6.55 | 6.31 | 0.08 | 0.094 | 0.021 | 0.603 |
| 90 | 6.18 | 6.09 | 6.32 | 6.08 | 0.10 | 0.477 | 0.113 | 0.487 |
| 120 | 6.20 | 5.96 | 5.96 | 5.75 | 0.11 | 0.049 | 0.052 | 0.853 |
| 180 | 5.98 | 5.89 | 5.72 | 5.62 | 0.13 | 0.058 | 0.476 | 0.980 |
| 300 | 6.03 | 5.90 | 5.70 | 5.71 | 0.83 | 0.009 | 0.479 | 0.434 |

1) $p < 0.05$: sign. differences between treatments

Table 4: Effects of feeding on the $\text{NH}_3\text{-N}$ -concentration (mmol/l) in rumen liquid before and after morning feeding

| Time after feeding (min) | Ration | | | | | Effects ¹⁾ | | |
|--------------------------|--------|--------|------|--------|------|-----------------------|-------------|-------|
| | H 70 | HLO 70 | H 30 | HLO 30 | PSEM | Hay | Linseed oil | HxLO |
| Before | 3.1 | 3.8 | 6.5 | 5.5 | 1.3 | 0.036 | 0.771 | 0.407 |
| 30 | 3.8 | 4.9 | 8.1 | 7.1 | 1.8 | 0.050 | 0.993 | 0.511 |
| 60 | 7.4 | 6.2 | 11.5 | 9.7 | 1.7 | 0.019 | 0.300 | 0.851 |
| 90 | 8.9 | 7.9 | 12.1 | 10.1 | 1.6 | 0.105 | 0.392 | 0.553 |
| 120 | 8.8 | 8.2 | 12.2 | 10.7 | 2.5 | 0.167 | 0.621 | 0.830 |
| 180 | 6.9 | 7.7 | 4.9 | 7.2 | 2.0 | 0.485 | 0.380 | 0.664 |
| 300 | 2.8 | 3.2 | 1.2 | 1.7 | 0.8 | 0.056 | 0.504 | 0.929 |

¹⁾ see Table 3

Low portions of hay in the ration (H 30 and HLO 30) reduced the rumen pH significantly at different times after feeding. These results are caused by a lower rumination and a more intensive fermentation of easily fermentable carbohydrates from concentrates, and are in agreement with some references (Flachowsky et al., 1993; Griinari et al., 1998; Kalscheur et al., 1997).

Except for the time 60 min. after the start of feeding, linseed oil did not significantly influence the rumen pH-

with concentrate rich rations (H 30, HLO 30). Later the $\text{NH}_3\text{-N}$ -concentration in rumen liquid was lower in the H 30 and HLO 30-rations in comparison with the rations rich in hay, probably caused by the high microbial activity and the N-demand during this time. Linseed oil did not significantly influence the $\text{NH}_3\text{-N}$ -concentration in rumen liquid (Table 4), which is in agreement with papers reviewed by Doreau and Ferlay (1995).

Apart from the valeric acid portion, the concentration

and pattern of short-chain fatty acids (SCFA) and the acetate: propionate ratio in the rumen liquid were not significantly influenced by ration composition and 200 g linseed oil supplementation (Table 5). Higher concentrations of acetic acid and lower values for propionic acid after feeding rations rich in roughages are described in many papers (e.g. Griinari et al., 1998; Ueda et al., 2003), but could be observed only in tendency

between individual measurements.

Nutrient flow and protein yield at the duodenum

The ration composition significantly influenced the apparent ruminal digestibility and the flow of organic matter (OM) and NDF to the duodenum (Table 6). The lower apparent ruminal digestibility of the OM of the

Table 5: Effect of feeding on the concentration of volatile fatty acids (VFA) in rumen liquid 180 min after morning feeding

| VFA | Ration | | | | PSEM | Effects ¹⁾ | | |
|-----------------------------|--------|--------|-------|--------|------|-----------------------|-------------|-------|
| | H 70 | HLO 70 | H 30 | HLO 30 | | Hay | Linseed oil | HxLO |
| Total VFA (mmol/l) | 144.6 | 140.4 | 136.5 | 149.6 | 13.1 | 0.967 | 0.738 | 0.519 |
| Molar proportions (%) | | | | | | | | |
| Acetic acid | 64.3 | 64.2 | 58.9 | 61.3 | 2.4 | 0.100 | 0.632 | 0.598 |
| Propionic acid | 19.9 | 19.5 | 20.2 | 20.4 | 1.6 | 0.724 | 0.923 | 0.896 |
| Butyric acid | 12.5 | 12.7 | 15.3 | 13.7 | 1.1 | 0.087 | 0.516 | 0.387 |
| Valeric acid | 1.7 | 1.8 | 3.6 | 3.0 | 0.3 | <0.001 | 0.426 | 0.363 |
| Acetic acid: Propionic acid | 3.3 | 3.4 | 3.2 | 3.3 | 0.4 | 0.801 | 0.857 | 0.967 |

¹⁾ see Table 3

Table 6: Nutrient flow to the duodenum

| Nutrients | Ration | | | | PSEM | Effects ¹⁾ | | |
|------------------------------|--------|--------|------|--------|------|-----------------------|-------------|-------|
| | H 70 | HLO 70 | H 30 | HLO 30 | | Hay | Linseed oil | HxLO |
| Flow (kg day ⁻¹) | | | | | | | | |
| OM | 5.50 | 5.70 | 5.85 | 6.05 | 0.14 | 0.019 | 0.167 | 0.986 |
| NDF | 2.22 | 2.32 | 1.99 | 1.83 | 0.10 | 0.001 | 0.749 | 0.199 |
| Ruminal digestibility (%) | | | | | | | | |
| OM | 55.0 | 53.0 | 52.0 | 50.6 | 1.0 | 0.042 | 0.335 | 0.888 |
| NDF | 65.6 | 63.7 | 54.3 | 58.3 | 1.8 | <0.001 | 0.582 | 0.132 |

¹⁾ see Table 3

($p > 0.05$, Table 5). Reasons for this could be the feeding frequency and the low level of DMI (see Table 2).

Some authors described a decrease in the acetic acid concentration in rumen liquid after supplementation of oil or other so-called rumen-unprotected fats (e.g. Doreau et al., 1993; Jenkins, 1993) because of the lower microbial fibre degradation. The present data did not show any significant influence of linseed oil on SCFA, probably because of the low amount of oil supplementation. Additionally some parameters of rumen fermentation failed to reach significance due to large differences

concentrate rich rations (H 30, HLO 30) was caused by more microbial OM at the duodenum (Table 8).

The amount of fermented organic matter was significantly higher with H 30 and HLO 30 as compared to H 70 and HLO 70 (Table 8).

The increased hay portion (H 70, HLO 70) in the ration and higher fibre intake (Table 2) resulted in an increased NDF-degradation in the rumen because of more stable pH but a higher NDF-flow to the duodenum as compared to the rations high in concentrate (H 30, HLO 30; Table 6).

The low amounts of supplemented linseed oil did not significantly influence ruminal digestibility and the duodenal flow of OM and NDF. The feeding of rations rich in concentrates (H 30, HLO 30) significantly increased the N-intake of cows, the N-flow to the duodenum, and the NAN at the duodenum (Table 7). The N-flow was significantly higher than the N-intake in all groups due to the microbial protein synthesis (see Table 8). Linseed oil supplementation showed no significant influence on parameters of N-flow to the duodenum (Table 7).

Higher amounts of unsaturated fatty acids have been under discussion for their ability to affect microbial protein

synthesis and efficiency of microbial protein synthesis, because of their depressant effects on protozoal populations (Doreau and Ferlay, 1995). Accordingly, the efficiency of microbial protein synthesis (g MP/kg FOM) tended ($p = 0.07$) to be higher with oil supplementation (Table 8).

The roughage/concentrate ratio significantly influenced the amounts of fermented organic matter (FOM) in the rumen, the microbial protein (MP) per day and per kg FOM, and the amounts of utilisable crude protein (nCP, Table 8).

The higher flow of NAN and nCP at the duodenum in animals fed with rations rich in concentrates

Table 7: Flow of Nitrogen to the duodenum

| Parameters | Ration | | | | | Effects ¹⁾ | | |
|---|--------|--------|-------|--------|------|-----------------------|-------------|-------|
| | H 70 | HLO 70 | H 30 | HLO 30 | PSEM | Hay | Linseed oil | HxLO |
| N-intake (g day ⁻¹) | 243.9 | 256.9 | 321.3 | 316.0 | 7.3 | <0.001 | 0.594 | 0.227 |
| N-Flow to the duodenum (g day ⁻¹) | 343.6 | 371.1 | 394.5 | 409.5 | 15.2 | <0.001 | 0.060 | 0.566 |
| (% of N-intake) | 141.1 | 145.0 | 123.2 | 125.9 | 5.2 | 0.005 | 0.312 | 0.785 |
| NAN ²⁾ at the duodenum (g day ⁻¹) | 325.6 | 351.8 | 373.9 | 389.3 | 10.3 | <0.001 | 0.056 | 0.604 |
| NAN/ME (g MJ ME ⁻¹) | 2.6 | 2.7 | 2.5 | 2.5 | 0.01 | 0.163 | 0.460 | 0.470 |

1) see Table 3

2) Non Ammonia-N

Table 8: Parameters of rumen fermentation and microbial protein yield

| Parameters | Ration | | | | | Effects ¹⁾ | | |
|---|--------|--------|--------|--------|------|-----------------------|-------------|-------|
| | H 70 | HLO 70 | H 30 | HLO 30 | PSEM | Hay | Linseed oil | HxLO |
| FOM ²⁾ (kg d ⁻¹) | 9.0 | 9.1 | 9.4 | 9.5 | 0.1 | 0.03 | 0.297 | 0.932 |
| (% OM intake) | 74.4 | 74.7 | 76.9 | 76.9 | 1.1 | 0.047 | 0.874 | 0.893 |
| MN ³⁾ (% of NAN) | 62.8 | 63.1 | 68.7 | 69.3 | 0.8 | <0.001 | 0.522 | 0.868 |
| MP ⁴⁾ (g d ⁻¹) | 1278.8 | 1387.5 | 1605.6 | 1688.1 | 45.8 | <0.001 | 0.048 | 0.782 |
| (g MJ ME ⁻¹) | 10.1 | 10.6 | 10.8 | 10.9 | 0.3 | 0.104 | 0.369 | 0.607 |
| (g kg FOM ⁻¹) | 142.2 | 152.6 | 171.3 | 177.7 | 4.4 | <0.001 | 0.071 | 0.654 |
| Endogenes protein ⁵⁾ (g d ⁻¹) | 157.4 | 165.3 | 160.6 | 166.5 | 3.6 | 0.551 | 0.072 | 0.784 |
| UDP ⁶⁾ (g d ⁻¹) | 599.1 | 646.4 | 571.2 | 578.4 | 28.6 | 0.108 | 0.351 | 0.491 |
| (% of CP-intake) | 27.9 | 27.8 | 23.2 | 22.6 | 0.8 | <0.001 | 0.659 | 0.801 |
| uCP (gd ⁻¹) | 1877.6 | 2033.6 | 2176.6 | 2266.7 | 62.8 | <0.001 | 0.062 | 0.605 |

¹⁾ see Table 3

²⁾ Fermented organic matter in the rumen = OM intake – (OM in the duodenum – OM of microorganisms); OM of microbes = 11.8 % microbial N

³⁾ Microbial N

⁴⁾ Microbial protein

⁵⁾ EP = 3.6 g/kg DM at the duodenum x 6.25

⁶⁾ UDP = uCP – MP ⁶⁾ uCP = (NAN x 6.25) – endogenes protein (EP)

bases on the increase of microbial protein synthesis (Table 8). The microbial protein synthesis per kg FOM varied between 142.2 and 177.7 g. The average of the H 70-rations amounted to 197.4, those of the H 30 – groups to 174.5 g per kg FOM. Kalscheur et al. (1997b) reduced the roughage portion from 60 to 25 % and measured an increase of MP from 195.0 to 316.3 g/kg FOM based on the reduction of FOM from 52 to 39 % of OM intake. The MP synthesis per kg FOM of the present study is in the range of measurements by Stern et al. (1994; \bar{x} : 181; range: 69 – 266 g MP/kg FOM) and Lebzién and Voigt (1999; \bar{x} : 188; range: 63 – 313 g MP/kg FOM). NRC (2001) considered an average of 186 g (75 – 338 g MP/kg FOM). There was no significant influence of linseed oil supplementation on parameters of N-flow and protein yield at the duodenum (Tables 7 and 8) in agreement with some references (Doreau and Ferlay, 1995; Kalscher et al., 1997a; Oldich and Firkins, 2000; Ueda et al., 2003).

REFERENCES

- CZERKAWSKI, J. W. – CLAPPERTON, J. L. 1984. Fats as energy-yielding compounds in the ruminant diet. In: *Fats in animal nutrition* by J. WISEMAN (ed.): Butterworths, 1984, p. 249-263.
- DE BRABANDER, D. L. – BE BOEVER, J. L. – DE SMET, A. M. – VANACKER, J. M. – BOU-CQUE, CH. V. – BOTTERMANN, S. M. 1999. Evaluation of physical structure in dairy cattle nutrition. In: *Recent Advances in Animal Nutrition* by P.C. Garnsworthy and J. Wiseman (Ed.) Nottingham: Univ. Press, 1999, p. 111-145.
- DIN 38406-E5-2. 1998. Deutsche Einheitsverfahren zur Wasser- Abwasser- und Schlammuntersuchung. Fachgruppe Wasserchemie in der GDChin Gemeinschaft mit dem Normenausschuss Wasserwesen im deutschen Institut für Normung e.V. 40th ed. Wiley-VCH Verlag GmbH, Wien: Weinheim und Beuth Verlag.
- DOREAU, M. – FERLAY, A. – ELMADDHA, Y. 1993. Organic matter and nitrogen digestion by dairy cows fed calcium salts of rapeseed oil fatty acids or rapeseed oil. In: *Journal Animal Science*, Vol. 71, 1993, p. 499-504.
- DOREAU, M. – FERLAY, A. 1995. Effect of dietary lipids on nitrogen metabolisms in the rumen: a review. In: *Livestock Production Science*, Vol. 43, 1995, p. 97-110.
- DROCHNER, W. – HELLER, R. 1995. Neue Ergebnisse geschützter Fette bei Wiederkäuern. In: *Übersichten Tierernährung*, Vol. 24, 1995, p.52-61.
- FLACHOWSKY, G. 1999. Fetteinsatz in der Milchkuhfütterung, Proc. In: *Aktuelle Themen der Tierernährung und Veredelungswirtschaft*. Cuxhaven: 27./28.10.1999, 1999, p. 83-105.
- FLACHOWSKY, G. – KOCH, H. – TIROKE, K. – MATTHE, M. 1993. Influence of the ratio between wheat straw and ground barley, ground corn and dried sugar beet pulp on in sacco dry matter degradation of ryegrass and wheat straw, rumen fermentation and apparent digestibility in sheep. In: *Archives Animal Nutrition*, Vol. 43, 1993, p. 157-167.
- FLACHOWSKY, G. – RAASCH, A. – RAASCH, R. v MÖCKEL, P. 1996. Influence of rumen protected fat with methionin on the performance of lactating dairy cows. In: *Journal Applied Animal Research*, Vol. 10, 1996, p. 135-148.
- GEISLER, C. – HOFFMANN, M. – HICKEL, B. 1976. Ein Beitrag zur gaschromatographischen Bestimmung flüchtiger Fettsäuren. In: *Archives Animal Nutrition*, Vol. 26, 1976, p. 123-129.
- GfE 2001. Energie- und Nährstoffbedarf landwirtschaftlicher Nutztiere, Nr. 8, Empfehlungen von Energie- und Nährstoffversorgung der Milchkühe und Aufzuchttrinder. DLG-Verlag, Frankfurt (Main), 2001, 136 p.
- GRIINARI, J. M. – DWYER DEBRA, A. – MCGUIRE, M. A. – BAUMAN, D. E. – PALMQUIST, D. L. – NURMELA, K. V. V. 1998. Trans-octadecenoic acids and milk fat depression in lactating dairy cows. In: *Journal Dairy Science*, Vol. 81, 1998, p. 1251-1261.
- GRIINARI, J.M. – BAUMAN, D.E. 2006. Milk fat depression concepts, mechanisms and management applications. In: K. Serjzen et al. (eds.), Ruminant physiology, Wageningen: Acad. Publ., p. 389-417.
- JAHREIS, G. – STEINHART, H. – PFALZGRAF, A. – FLACHOWSKY, G. – SCHÖNE, F. 1996. Zur Wirkung von Rapsölfütterung an Milchkühe auf das Fettsäurenspektrum des Butterfettes. *Z. Ernährungswissenschaft*, Vol. 35, 1996, p. 185-190.
- JENKINS, T.C. 1993. Symposium: advances in ruminant lipid metabolism – Lipid metabolism in the rumen. In: *Journal Dairy Science*, Vol. 76, 1993, p. 3851-3863.
- JILG, T. – AIPLE, A. P. – STEINGASS, M. 1988. Fettstoffwechsel und Wirkungen von Futterfetten beim Wiederkäuer. In: *Übersichten Tierernährung*, Vol. 16, 1988, p. 109-152.
- KALSCHOUR, K.F. – TETER, B.B. – PIPEROVA, L.S. – ERDMAN, R.A. 1997a. Effect of fat source on duodenal flow at trans-C_{18:1} fatty acids and milk fat production in dairy cows. In: *Journal Dairy Science*, Vol. 80, 1997, p. 2115-2126.
- KALSCHOUR, K. F. – TETER, B. B. – PIPEROVA, L. S. – ERDMAN, R. A. 1997b. Effect of dietary forage concentration and buffer addition on duodenal flow of trans-C_{18:1} fatty acids and milk fat production in dairy cows. In: *Journal Dairy Science*, Vol. 80, 1997, p. 2104-2114.
- KENNELLY, J. J. – GLIMM, D. R. 1998. The biological potential to alter the composition of milk. In: *Canadian Journal Animal Science*, Vol. 78, Suppl., 1998, p. 23-56.
- LEBZIEN, P. – PAUL, C. 1997. Use of near-infrared reflectance spectroscopy for the estimation of the microbial portion of non-ammonia-nitrogen in the duodenum of dairy cows. In: *Animal Feed Science Technology*, Vol. 68, 1997, p. 225-233.
- LEBZIEN, P. – VOIGT, J. 1999. Calculation of utilizable crude protein at the duodenum of cattle by two different approaches. In: *Archives Animal Nutrition*, Vol. 52, 1996, p. 1363-1368.
- MACHMÜLLER, A. – KREUZER, M. 1999. Methane suppression by coconut oil and associated effects on nutrient and energy balance in sheep. In: *Canadian Journal Animal Science*, Vol. 79, 1999, p. 65-72.
- MEYER, U. – GÄDEKEN, D. – SPOLDERS, M. – FLACHOWSKY, G. 2001. Physical structure value systems in dairy cattle feeding. In: *Proceedings Society Nutrition Physiology*, Vol. 10, 2001, 141 p.

- MEYER, U. – GÄDEKEN, D. – SPOLDERS, M. – FLACHOWSKY, G. 2002. A comparison of recommendations for the fibre supply of dairy cows fed Total Mixed Rations (TMR). In: *Proceedings Society Nutrition Physiology*, Vol. 11, 2002, 133 p.
- NAUMANN, C. – BASSLER, R. 1993. Die chemische Untersuchung von Futtermitteln. Darmstadt: VDLUFA-Verlag, 1993.
- NRC-NUTRIENT REQUIREMENTS OF DAIRY CATTLE. 2001. Washington D.C.: 7th Ed. National Academy press., 381 p.
- OLDICK, B.S. – FIRKINS, J.L. 2000. Effects of degree of fat saturation on fiber digestion and microbial protein synthesis when diets are fed twelve times daily. In: *Journal Animal Science*, Vol. 78, 2000, p. 2412-2420.
- PALMQUIST, D. L. – JENKINS, T. C. 1980. Fat in lactation rations: Review. In: *Journal Dairy Science*, Vol. 63, 1980, p. 1-15.
- PIATKOWSKI, B. – NAGEL, S. – BERGNER, E. 1977. Das Wiederkauverhalten von Kühen bei unterschiedlicher Trockensubstanzaufnahme und verschiedener physikalischer Form von Grasheu. In: *Archiv Tierernährung*, Vol. 27, 1977, p. 563-569.
- ROHR, K. – BRANDT, M. – LEBZIEN, P. – SCHAFFT, H. 1984. Measurement of duodenal flow in dairy cows by either total collection or spot sampling, using a special cannula. In: *Canadian Journal Animal Science*, Suppl., 1984, p. 116-117.
- ROHR, K. – BRANDT, M. – CASTRILLO, O. – LEBZIEN, P. – ASSMUS, G. 1979. Der Einfluss eines teilweisen Ersatzes von Futterprotein durch Harnstoff auf den Stickstoff- und Aminosäureneinfluss am Duodenum. In: *Landbauforschung Völkenrode*, Vol. 29, 1979, p. 32-40.
- STERN, M. D. – VAGRA, G. A. – CLARK, J. H. – FIRKINS, J. L. – HUBER, J. T. – PALMQUIST, D. L. 1979. Evaluation of chemical and physical properties of feeds that affect protein metabolism in the rumen. In: *Journal Dairy Science*, Vol. 76, 1979, p. 2762-2786.
- SUTTON, J. D. – KNIGHT, R. – MC ALLAN, A. B. – SMITH, R. H. 1983. Digestion and synthesis in the rumen of sheep given diets with free and protected oils. In: *British Journal Nutrition*. Vol. 49, 1983, p. 419-432.
- UEDA, K. – FEERLAY, A. – CHABROT, J. – LOOR, J.J. – CHILLIARD, Y. – DOREAU, M. 2003. Effect of linseed oils supplementation on ruminal digestion in dairy cows fed diets with different forage: concentrate ratios. In: *Journal Dairy Science*, Vol. 86, 2003, p. 3999-4007.
- WAGNER, K. – AULRICH, K. – LEBZIEN, P. – FLACHOWSKY, G. 1998. Research Note: Effect of duodenal-infused unsaturated fatty acids on dairy milk composition. In: *Archives Animal Nutrition*, Vol. 51, 1998 p. 349-354.
- WILLIAMS, C.H. – DAVID, D.J. – IISMAA, O. 1962. The determination of chromic oxide in faeces samples by atomic absorption spectrometry. In: *Journal Agriculture Science*, Vol. 59, 1962, p. 381-385.

Authors' address: Prof. Dr. Gerhard Flachowsky, Institute of Animal Nutrition of the Federal Agricultural Research Centre, Bundesallee 50, 38116 Braunschweig, Germany