

Scientific paper formerly presented

EVALUATION OF SELECTED BIOCHEMICAL PARAMETERS IN BLOOD PLASMA, URINE AND MILK OF DAIRY COWS DURING THE LACTATION PERIOD

T. FILIPEJOVÁ*, J. KOVÁČIK

Slovak University of Agriculture, Nitra, Slovak Republic

ABSTRACT

The aim of our study was to determine selected biochemical parameters in the blood, urine and milk of dairy cows. Fifteen dairy cows from selected agricultural farms were divided into three groups as follow: group I- 3-4 weeks after calving (the beginning of lactation), group II - 3-4 months after calving (the middle of lactation), and group III - 2-3 weeks before calving (the dry period). Concentrations of selected biochemical parameters in the blood (aspartate aminotransferase – AST, urea, total proteins - TP, glucose, and cholesterol), urine (calcium - Ca, phosphorus - P, urea and pH), and milk (fats, proteins, lactose, solids, Somatic Cell Count, urea, milk freezing point and rennetability) were measured. Glucose concentration (3.22 ± 0.21 mmol.l⁻¹) was significantly lower at the beginning of lactation ($p < 0.05$) in comparison with the middle of lactation (3.69 ± 0.08 mmol.l⁻¹) and the dry period (3.74 ± 0.21 mmol.l⁻¹). However, concentration of AST (1.42 ± 0.31 μkat.l⁻¹) was significantly higher ($p < 0.01$) at the beginning of lactation than in the dry period (0.88 ± 0.31 μkat.l⁻¹). Concentration of cholesterol (6.22 ± 0.76 mmol.l⁻¹) in the middle of lactation was significantly higher ($p < 0.001$) than those at the beginning of lactation (3.82 ± 0.67 mmol.l⁻¹) and dry period (2.842 ± 0.67 mmol.l⁻¹). Cows in the middle of lactation had a higher freezing point of milk (526.20 ± 1.92 -m.°C) ($p < 0.01$) in comparison with dairy cows at the beginning of lactation (520.00 ± 2.74 -m.°C). Some other symptoms leading to sub-clinical diseases, besides worsening the technological quality of the cow milk, were found.

Key words: biochemical parameters; metabolism; metabolic disorders; milk quality

INTRODUCTION

Various factors influence milk quantity and quality. The quality of milk and its nutritional value influenced by many factors (Tančin et al., 2006; Tančin et al., 2007). One of them is the milking process itself, in which the emphasis is to minimize the undesirable contaminant microflora, which significantly offset the technological quality of milk and may cause health injury (Kirchnerová and Foltys, 2005). Specifically, the nutrition and health status are major factors in determining cow milk characteristics. Metabolic perturbations, lack of nutrition and sub-clinical disorders could be detected by measuring some metabolic parameters in the blood, urine and milk (Slanina and Beseda, 1992). Such parameters are known to influence the composition and technological

aspects of milk quality in dairy cows (Illek et al., 2002; Kováčik et al., 2004). Several factors like nutrition, metabolic status, production, health and the management affect dairy cow's fertility (Roxström et al., 2000; Clark et al., 2000; Mwaanga and Janowski, 2000; Mihm, 1999; Kruij et al., 1998). The changes that take place in the endocrine system of cows in the dry period and still more at early lactation intensify gluconeogenesis, lipolysis and ketogenesis (Ling et al., 2003).

Dairy cattle, like most lactating mammals, are usually in negative energy balance within the first few weeks of lactation (Nielsen, 1999). After the delivery, the cow requires high amount of energy from body reserves, but actually it is not able to cover the required nutrients consumed, what leads to the loss of body weight. The continuous increase in milk production has created new

*Correspondence: E-mail: filipejova@gmail.com

Ing. Terézia Filipejová, Slovak University of Agriculture, Faculty of Biotechnology and Food Sciences,
Department of Animal Physiology, Tr. A. Hlinku 2, 949 76 Nitra, Slovak Republic;

Tel: +410 37 641 4288

challenges for high-producing dairy cows, especially during the transition period. At the beginning of lactation, dairy cows have to cope with the high energy and protein demands for milk synthesis at the time when nutrient intake is low. Mobilizing energy and protein from body tissue stores and repartition of nutrients away from extra-mammary tissues are the primary alternatives to supply sufficient nutrients for milk production during the first weeks of lactation. Excessive body reserves, especially fat, can cause a series of metabolic disorders and consequent production losses (Fourichon et al., 1999).

Furthermore, physiological and pathological changes associated with negative energy balance are important factors related to development of ketosis, displaced abomasum, and retained placenta (Duffield *et al.*, 2002).

The aim of our study was a determination of selected biochemical parameters in the blood, urine and milk of dairy cows and their analysis.

MATERIAL AND METHODS

Fifteen dairy cows from selected agricultural farms were divided into three groups: group I - 3-4 weeks after calving (the beginning of lactation), group II - 3-4 months after calving (the middle of lactation) and group III - 2-3 weeks before calving (the dry period).

Blood samples for biochemical analysis were taken from *vena jugularis* 2 hours after morning feeding. The blood plasma was separated from whole blood by the centrifugation at 3000 rpm for 30 minutes and samples

were stored at $-18\text{ }^{\circ}\text{C}$. Selected biochemical parameters in blood plasma (aspartate aminotransferase (AST), urea, total proteins (TP), glucose (GLU) and cholesterol (CHOL), in urine (calcium (Ca), phosphorus (P), urea, pH) were analyzed using semi-automated clinical chemistry analyzer Microlab 300 Vilat Scientific, Dieren, The Netherlands) (Massanyi et al., 1995; Massanyi et al., 2007).

Samples of milk were cooled down until $6\text{ }^{\circ}\text{C}$ was reached. Samples were kept at the same temperature during the determination of milk quality parameters: Somatic Cell Counts (by the Fossomatic device), urea (photocolorimetric device with Ehrlich's reagent, 530 nm wavelength), Non-Fat-Solis (by the MilkoScan apparatus), acidity, rennetability, fermentace ability and thermostability. Significant differences between groups of dairy cows were evaluated by the Scheffe's - test. Differences between the groups at $p<0.05$, $p<0.01$ or $p<0.001$ were considered as significant.

RESULTS AND DISCUSSION

Blood biochemical parameters of animals are affected by seasonal variations (Massányi et al., 2009) or food additives (Arpášová et al., 2009). Measurements of blood metabolites are associated with great variations, indicating large fluctuations in metabolite profiles during the peri-parturient period (Drackley, 1999).

Concentrations of biochemical parameters in blood serum, urine and milk during lactation period are shown in tables 1-3.

Table 1: Concentration of AST, urea, total proteins, glucose, cholesterol in blood plasma of dairy cows at the beginning of lactation, in the middle of lactation and the dry period

	Variable	Minimum	Maximum	x	s	cv
BL	AST ($\mu\text{kat.l}^{-1}$)	1.15	1.90	1.42**	0.31	21.85
	Urea(mmol.l^{-1})	2.18	3.45	2.94	0.48	16.20
	TP (g.l)	72.00	88.00	79.00	6.44	8.15
	GLU(mmol.l^{-1})	2.87	3.45	3.22*	0.21	6.64
	CHOL(mmol.l^{-1})	2.90	4.60	3.82	0.67	17.50
ML	AST ($\mu\text{kat/L}$)	1.04	1.53	1.28	0.18	14.18
	Urea(mmol.l^{-1})	2.87	5.53	4.82	1.17	24.17
	TP(g.l)	73.00	90.00	78.80	7.26	9.21
	GLU(mmol.l^{-1})	3.61	3.82	3.69	0.08	2.19
	CHOL(mmol.l^{-1})	5.10	7.00	6.22***	0.76	12.21
D	AST ($\mu\text{kat.l}$)	0.77	1.05	0.88	0.31	21.85
	Urea(mmol.l^{-1})	0.19	0.40	0.30	0.08	25.50
	TP(g.l)	56.00	85.00	73.60	6.44	8.15
	GLU(mmol.l^{-1})	3.43	4.39	3.74	0.21	6.64
	CHOL(mmol.l^{-1})	2.00	3.90	2.84	0.67	17.50

BL-beginning of lactation, ML-middle of lactation, D-dry period, AST- aspartate aminotransferase, TP-total proteins, GLU-glucose, CHOL-cholesterol, x -mean, s-standard deviation, cv- coefficient of variation, * $p<0.05$, ** $p<0.01$, *** $p<0.001$

Table 2: Concentration of calcium, phosphorus, urea and pH in urine of dairy cows at the beginning of lactation, in the middle of lactation and the dry period

	Variable	Minimum	Maximum	x	s	cv
BL	Ca (mmol.l ⁻¹)	0.11	4.39	2.52	2.06	82.03
	P (mmol.l ⁻¹)	0.09	2.37	0.88	1.05	118.97
	Urea(mmol.l ⁻¹)	72.00	372.00	223.25	126.45	56.64
	pH (logmole)	8.00	8.30	8.13	0.15	1.85
ML	Ca (mmol.l ⁻¹)	0.22	1.17	0.70	0.36	51.35
	P (mmol.l ⁻¹)	0.10	8.38	1.84	3.66	198.54
	Urea(mmol.l ⁻¹)	61.00	245.00	179.20	75.05	41.88
	pH (logmole)	7.80	8.30	8.12	0.20	2.52
D	Ca (mmol.l ⁻¹)	0.96	3.50	1.85	1.13	61.32
	P (mmol.l ⁻¹)	0.09	0.86	0.35	0.35	101.24
	Urea(mmol.l ⁻¹)	135.00	468.00	293.25	136.76	46.64
	pH (logmole)	8.00	8.30	8.15	0.13	1.58

BL-beginning of lactation, ML-middle lactation, D-dry period, Ca-calcium, P-phosphorus, x-mean, s-standard of deviation, cv- coefficient of variation

Pregnancy and lactation have a great impact on the intensity of metabolism and on metabolic parameters in the blood (Tab. 1). The activity of blood aminotransferases is very important because they act as a catalyst in connection to the metabolism of amino acids and carbohydrates. Changes in their activity in the blood can be a consequence of their increased activity in cells, or damages in cell structure (Milinković-Tur, 2005). The concentration of AST ($1.42 \pm 0.31 \mu\text{kat.l}^{-1}$) in dairy cows at the beginning of lactation was significantly

higher ($p < 0.01$) in comparison with the dry period ($0.88 \pm 0.31 \mu\text{kat.l}^{-1}$). AST is a widely distributed enzyme, which is found in many tissues and organs, with high activity in the liver (Zimmerman et al., 1968). Increased AST activity in the serum is a sensitive marker of liver damage (Meyer and Harvey, 1998). The intensity of metabolic changes, mainly of proteins, is also reflected in the concentration of other biochemical indicators of the blood, such as total protein, urea or aspartate aminotransferase (Falkenberg et al., 1997; Kapelański

Table 3: Concentration of fats, proteins, lactose, somatic cell counts, urea, freezing point, rennetability, in milk of dairy cows at the beginning of lactation, and in the middle of lactation

	Variable	Minimum	Maximum	x	s	cv
BL	Fats (g.100g ⁻¹)	0.57	1.94	1.20	0.59	4950
	Proteins (g.100g ⁻¹)	2.93	3.58	3.17	0.26	8.07
	Lactose (g.100g ⁻¹)	4.73	5.01	4.93	0.12	2.47
	Solids (g.100g ⁻¹)	9.16	10.79	9.99	0.71	7.14
	Somatic Cell Counts (tis . ml ⁻¹)	9.00	366.00	134.20	15466	115.25
	Urea (mmol.l ⁻¹)	18.91	33.84	26.05	7.03	27.00
	Freezing point (-m. °C)	517.00	524.00	520.00	274	0.53
	Rennetability(s)	17.40	26.25	20.63	3.52	17.07
ML	Fats (g.100g ⁻¹)	0.66	3.27	1.61	1.20	74.34
	Proteins (g.100g ⁻¹)	2.69	3.85	3.05	0.47	15.26
	Lactose (g.100g ⁻¹)	4.02	4.99	4.59	0.44	56.64
	Solids (g.100g ⁻¹)	8.64	11.64	9.97	1.32	13.27
	Somatic Cell Counts (tis . ml ⁻¹)	20.00	6136.00	1525.40	2619.89	171.75
	Urea (mmol.l ⁻¹)	20.16	35.23	26.83	6.55	24.43
	Freezing point (-m. °C)	523.00	528.00	526.20**	1.92	0.37
	Rennetability(s)	20.62	46.12	34.17	11.79	34.50

BL-beginning of lactation, ML-middle of lactation, x - mean, s - standard of deviation, cv - coefficient of variation

** $p < 0.01$

et al., 2000; Więcek and Skomial, 2000). Measurements of blood metabolites require more frequent sampling to capture the dynamic changes in the peri-parturient period. The concentration of cholesterol (6.22 ± 0.76 mmol.l⁻¹) in the middle of lactation was significantly higher ($p < 0.001$) in comparison with concentration at the beginning of lactation (3.82 ± 0.67 mmol.l⁻¹) and dry period (2.842 ± 0.67 mmol.l⁻¹). Cholesterol increase in the course of lactation together with intensive steroid synthesis has been reported in several studies (Bösö et al., 2000; Pysera and Opalka, 2000).

The concentration of glucose (3.22 ± 0.21 mmol.l⁻¹) in dairy cows at the start of lactation was significantly lower ($p < 0.05$) in comparison with those in the middle of lactation (3.69 ± 0.08 mmol.l⁻¹) and at the dry period (3.74 ± 0.21 mmol.l⁻¹).

We did not observe any significant differences in urine indicators of dairy cows (Tab. 2).

Studies on influences of the milk yield increase on cow's health status and milk quality becoming more and more important (Sawa and Piwczynski, 2002; Ayadi et al., 2003; Remond et al., 2004; Junqueir, et al., 2005). However, it is possible only for a limited number of milk indicators, such as the milk yield, fat, protein, lactose and urea contents and somatic cell counts. The milk yield is an important economic and health factor closely connected with the health status of dairy cows, their reproduction performance, longevity and milk composition and properties (Janů et al., 2007). We found, that freezing point of milk (526.20 ± 1.92 -m.°C) in the middle of lactation was significantly different ($p < 0.01$) compared to dairy cows at the beginning of lactation (520.00 ± 2.74 -m.°C) (Tab. 3).

CONCLUSION

In conclusion, deficiency in dairy cow's nutrition may influence many biochemical and physiological processes. Consequently they disturb the relationship between metabolic capacities of animals required for milk production and, thereby, cause metabolic disorders.

In this study some significant differences related to the lactation period were found. Glucose concentration at the beginning of lactation was significantly lower ($p < 0.05$) in comparison with the middle of lactation and the dry period. Concentration of AST at the beginning of lactation was significantly higher ($p < 0.01$) in comparison with the dry period. Concentration of cholesterol in the middle of lactation was significantly higher ($p < 0.001$) in comparison with the beginning of lactation and the dry period. Freezing point of milk in the middle of lactation was significantly higher ($p < 0.01$) in comparison to dairy cows at the beginning of lactation. Obtained results could serve to better understanding of biochemical processes in

dairy cows for estimating their physiological status.

REFERENCES

- AYADI, M. – CAJA, G. – SUCH, X. – KNIGHT, CH. 2003. Effect of omitting one milking weekly on lactation performances and morphological udder changes in dairy cows. *Journal of Dairy Science*, 2003, vol. 86, p. 2352-2358.
- ARPASOVA, H. – ANGELOVICOVA, M. – HASCÍK, P. – CAPCAROVA, M. – KOLESAROVA, A. – KACANIOVA, M. – HANOVA, M. 2009. Influence of plant essential oils on selected qualitative parameters of hen table eggs. *Acta fytotechnica and zootechnica*, 2009, no. 12, p. 8-15.
- BÖSÖ, A. R. – SAUKKO, T. M. – TESFA, A. T. – LINDBERG, L. A. 2000. Fat infiltration in liver and activity of lecithin: cholesterol acyltransferase in serum of dry and lactating dairy cows. *Research in Veterinary Science*, vol. 68, 2000, p. 169-173.
- CLARK, B. A. – CHAGAS, P. M. – GORE, B. – VERKERK, G. A. 2000. Prediction of post-partum anovulatory interval in dairy cows. – Proceedings of the New Zealand Society of Animal Production, vol. 60, 2000, p. 15-18.
- DRACKLEY, J. K. 1999. Biology of dairy cows during the transition period: The final frontier? *Journal of Dairy Science*, vol. 82, 1999, no. 11, p. 2259-2273.
- DUFFIELD, T. – BAGG, R. – DESCOTEAUX, L. – BOUCHARD, E. – BRODEUR, M. – DUTREMBLAY, D. – KEEFE, G. – LEBLANC, S. – DICK, P. 2002. Parturition monensin for the reduction of energy associated disease in postpartum dairy cows. *Journal of Dairy Science*, vol. 85, 2002, p. 397-405.
- FALKENBER, G. H. – RENNE, U. – LANGHAMMER, M. 1997. Effects of long-term selection on variation of blood metabolic substances in different mouse lines. Proceedings of the 48th Annual Meeting of the EAAP, 1997. Vienna, 25-28 August Book of Abstracts, p. 41.
- FOURICHON, C. H. – SEEGER, BAREILLE, N. – BEAUDEAU, F. 1999. Effects of disease on milk production in dairy cow: A review. *Preventive Veterinary Medicine*, vol. 41, p. 1-35.
- ILLEK, J. 2002. Vliv výživy na zdravotní stav dojníc a kvalitu mléka [Influence of nutrition on health status of dairy cows and milk quality]. *Výživa a zdraví vysokoprodukčních dojníc*. Brno : Veteduca, spol. s.r.o., 2002, s.7-9.
- JANŮ, L. – HANUŠ, O. – FRELICH, J. – MACEK, A. – ZAJÍČKOVÁ, I. – GENČUROVÁ, V. – JEDELSKÁ, R. 2007. Influences of Different Milk Yields of Holstein Cows on Milk Quality Indicators in the Czech Republic. *Acta Veterinaria Brno*, vol. 76, 2007, p. 553-561.
- JUNQUEIR, F. S. – MADALE, F. E. – REIS, G. L. 2005. Production and economic comparison of milking F-1 Holstein × Gir cows with and without the stimulus of the calf. *Livestock Production Science*, vol. 97, 2005, p. 241-252.
- KAPELANSKI, W. – PODKÓWKA, Z. – GRAJEWSKA, S. 2000. Poziom niektórych metabolitów surowicy krwi a efektywność tuczu (Level of certain blood serum metabolites and fattening efficiency in pigs). In Polish, with English summary. *Zeszyty Naukowe Przeglądu Hodowlanego*, vol. 48, 2000, p. 153-158.

- KIRCHNEROVA, K. – FOLTYS, V. 2005. Nové trendy v hodnotení kontaminujúcej mikroflóry v procese získavania a spracovania mlieka. [New trends in the evaluation of contaminant microflora in the process of obtaining and processing of milk] *Agriculture*, vol. 51, 2005, p. 308-314.
- KOVACIK, J. – KRAMAROVA, M. – MASSANYI, P. – FABIS, M. – BUKOVINSKY, M. 2004. Content of urea in biological fluids of dairy cows and technological quality of milk. *Risk factors of food chain*, Nitra, 2004, p. 128-130.
- KRUIP, T. A. M. – MEIJER, G. A. L. – RUKKWAMSUK, T. – WENSING, T. 1998. Effects of feeding in the dry period on fertility of dairy cows post partum. *Reproduction in Domestic Animals*, vol. 33, 1998, p. 165-168.
- LING, K. – JAAKSON, H. – SAMARÜTEL, J. – LEESMÄE, A. 2003. Blood metabolites of Estonian Holstein cows and their relation to some fertility parameters. *Agraarteadus*, vol. 14, 2003, no. 5, p. 269-276.
- MASSANYI, P. – TOMAN, R. – VALENT, M. – JONES, Q. 1995. Serum mineral profile of rabbits after an experimental administration of cadmium. *Journal of Environmental Science and Health*, 1995, p. 2221-2227.
- MASSANYI, P. – KOVACIK, J. – TOMAN, R. – LUKAC, N. – STAWARZ, R. – CAPCAROVA, M. – KOLESAROVA, A. 2007. Concentration of selected serum parameters after cadmium administration. *Physiological Research*, vol. 56, 2007, p. 22.
- MASSANYI, P. – SLAMECKA, J. – LUKAC, N. – CAPCAROVA, M. – MERTIN, D. – JURCIK, R. 2009. Seasonal variations in the blood biochemistry of brown hare. *Medycyna Weterynaryjna.*, vol. 65, 2009, no. 6, p. 389-393.
- MWAANGA, E. S. – JANOWSKI, T. 2000. Anestrus in dairy cows: causes, prevalence and clinical signs. – Review article. *Reproduction in Domestic Animals*, vol. 35, 2000, p. 193-200.
- MIHM, M. 1999. Delayed resumption of cyclicity in postpartum dairy and beef cows. *Reproduction in Domestic Animals*, vol. 34, 1999, p. 277-284.
- MILINKOVIĆ-TUR, S. – PERIĆ, V. – STOJEVIĆ, Z. – ZDELAR-TUK, M. – PIRSLJIN, J. 2005. Concentrations of total proteins and albumins, and AST, ALT and GGT activities in the blood plasma of mares during pregnancy and early lactation. *Veterinarski arhiv*, vol. 75, 2005, no.3, p.195-202.
- MEYER, D. J. – HARVEY, J. W. 1998. Veterinary Laboratory Medicine. Interpretation and Diagnosis. 2nd ed., W. B. Saunders Company : Philadelphia, London, Toronto, Montreal, Sydney, Tokyo. 1999, p. 157-187.
- NIELSEN, B. L. 1999. Perceived welfare issues in dairy cattle, with special emphasis on metabolic stress. In: *Metabolic Stress in Dairy Cows*. Oldham, J. D. - Simm, G. - Groen, A. F. - Nielsen, B. L. - Pryce, J. E. - Lawrence, T.L.J. In: *British Society of Animal Science, Occasional Publication*, vol. 24, 1999, p. 1-8
- PYSERA, B. – OPALKA, A. 2000. The Effect of Gestation and Lactation of Dairy Cows on Lipid and Lipoprotein Patterns and Composition in Serum During Winter and Summer Feeding. *Journal of Animal and Feed Science*, vol. 9, 2000, p. 411-424.
- REMOND, B. – POMIES, D. – DUPONT, D. – CHILIARD, Y. 2004. Once-a-day milking of multiparous Holstein cows throughout the entire lactation: milk yield and composition, and nutritional status. *Animal Research*, vol. 53, 2004, p. 201-212.
- ROXTRÖM, A. – STRANDBERG, E. – BERGLUND, B. – EMANUELSON, U. – PHILIPSSON, J. 2000. Genetic and environmental correlations among female fertility traits and milk production in different parities of Swedish Red and White Dairy Cattle. – Acta Agraria Scandinavia, Sect A, In: *Animal Science*, vol. 51, 2000, p. 7-14.
- SAWA, A. – PIWCZYNSKI, D. 2002. Somatic cell count and milk yield and composition in White × Holstein-Friesian cows. *Medycyna Weterynaryjna*, vol. 8, 2002, no. 28, p.636-640.
- SLANINA, L. – BESEDA, I. 1992. *Metabolický profil hovädzieho dobytku vo vzťahu k zdraviu a produkcii*. [Metabolic profile test of dairy cows in relationship of health and production] 2 vyd. Bratislava : ŠVS SR. 1992, s. 89. ISBN 80-71-48-001-0.
- TANCIN, V. – KIRCHNEROVA, K. – FOLTYS, V. – MACUHOVA, L. – TANCINOVA, D. 2006. Microbiological contamination and somatic cell count of bovine milk stripped before and after udder preparation for milking. *Slovak J. Anim. Sci.*, vol. 39, 2006, no.4, p. 214 – 217.
- TANCIN, V. – UHRINACAT, M. – MIHINA, S. – SUDZINOVA, J. – FOLTYS, V. – TANCINOVA, D. 2007. Somatic cell count and quarter milk flow parameters from udder of dairy cows. *Slovak J. Anim. Sci.*, vol. 40, 2007, no. 2, p. 79-82.
- WIĘCEK J. – SKOMIAŁ J., 2000. The effect protein and amino acids on urea level in blood of fattening pigs. Proceedings of the 51st Annual Meeting of the EAAP, 21-24 August, the Hague. 2000, p. 348.
- ZIMMERMAN, H. J. – DUJOVNE, C. A. – LEVY, R. 1968. The correlation of serum levels of two transaminases with tissue levels in six vertebrate species. *Comparative Biochemistry and Physiology*, vol. 25, p. 1081-1089.