

CELL RECEPTORS MEDIATING COMMUNICATION BETWEEN PREIMPLANTATION EMBRYO AND SURROUNDING ENVIRONMENT: CLUES FROM MOUSE AND RABBIT MODELS: A REVIEW

Štefan ČIKOŠ*, Alexandra ŠPIRKOVÁ, Janka KUBANDOVÁ, Veronika KOVAŘÍKOVÁ, Zuzana ŠEFČÍKOVÁ, Dušan FABIAN, Juraj KOPPEL

Centre of Biosciences of the Slovak Academy of Sciences, Institute of Animal Physiology, Košice, Slovakia

ABSTRACT

Preimplantation period of embryo development is one of the most sensitive phases in the mammalian ontogeny, and disturbances at this developmental stage can result in poor pregnancy outcomes (in both embryos resulting from natural conception or from biotechnology procedures). Results of experimental studies have shown that maternal physiological conditions and external environmental factors can significantly influence preimplantation embryo development, indicating a communication between the early embryo and its environment. The study of communication between the early embryo and surrounding environment has been focused mainly on protein signaling molecules, such as growth factors and cytokines. However, small-molecule ligands, such as biogenic monoamines, have been shown to influence preimplantation embryo development as well, and results obtained on mouse and rabbit models indicate that biogenic monoamine receptors are expressed in preimplantation embryos. Several adrenergic, dopamine, serotonin and histamine receptors were detected in mouse and rabbit ovulated oocytes and preimplantation embryos, and in mouse embryonic stem cells. Although the physiological role of biogenic monoamine receptors in early embryonic cells is not fully understood, experimental data indicate their involvement in the regulation of cell proliferation, differentiation and survival under physiological as well as unfavorable or pathological conditions (e.g. during maternal stress).

Key words: preimplantation embryo; cell receptors; embryo-maternal communication

INTRODUCTION

An important part of animal biotechnology is focused on animal reproduction, and number of approaches, such as artificial insemination, multiple ovulations, *in vitro* fertilization, embryo culture, embryo transfer, cryopreservation of gametes and embryos, nuclear transfer or cloning, transgenesis and embryonic stem cell production, are used in this field. Fertilized oocyte (zygote), which can result from natural conception or artificial insemination, develops several days in the oviduct, and finally it implants into the uterine wall

at the blastocyst stage. In some assisted reproductive technologies, the preimplantation development takes place *in vitro* and the blastocyst is then transferred into the uterus. Alternatively, the inner cell mass of blastocysts can serve as a source of embryonic stem cells, which are potentially usable in regenerative medicine.

Although the preimplantation period of development lasts only for a relatively short time, it represents one of the most sensitive phases in mammalian ontogeny (up to 50 % of embryo loss occurs during this period), and disturbances at this developmental stage can result in poor

*Correspondence: E-mail: cikos@saske.sk
Štefan Čikoš, Centre of Biosciences of the Slovak Academy
of Sciences, Institute of Animal Physiology,
Šoltésovej 4-6, 04001, Košice, Slovakia

Received: November 20, 2018

Accepted: December 3, 2018

pregnancy outcomes in both humans and animals (Nepomnaschy *et al.*, 2006; Humblot, 2001; Diskin *et al.*, 2008). The preimplantation embryo can develop relatively autonomously in very simple culture media, utilizing transcripts and proteins provided by the oocyte and gene products synthesized by an embryo itself (after activation of the embryonic genome). However, there are data indicating that *in vitro* culture of preimplantation embryos can alter gene expression and epigenetic reprogramming, and can lead to abnormal fetal and perinatal growth as well as to physiological and behavioral alterations in adulthood (for review see Khosla *et al.*, 2001; Fleming *et al.*, 2004; Ventura-Junca *et al.*, 2015). Moreover, results of animal studies and observations in humans have shown that maternal physiological condition (nutritional status, body condition, metabolic and other disorders, stress) as well as external environmental factors, e.g. various xenobiotics that can contaminate food, can significantly influence preimplantation embryo development (Kwong *et al.*, 2000; von Borell *et al.*, 2007; Fabian *et al.*, 2010; Kubandová *et al.*, 2014; Burkuš *et al.*, 2013; 2015; Janštová *et al.*, 2017; Babel'ová *et al.*, 2017). These data indicate that communication between the embryo and its environment takes place already in very early developmental stages.

The study of communication between the early embryo and surrounding environment has been mainly focused on protein signaling molecules, such as growth factors and cytokines, and relatively many receptors for protein ligands have been identified in preimplantation embryos (for review see Hardy and Spanos, 2002; Thouas *et al.*, 2015). However, we and others have demonstrated that small-molecule ligands, such as biogenic monoamines, can significantly influence preimplantation embryo development as well. This minireview focuses on the expression of receptors for biogenic monoamines and their potential role in early embryo development.

Laboratory mouse is the most widely used model for the study of early mammalian embryo development and most information on mechanisms involved in embryo-maternal communication comes from the mouse model. Recently, the progress in rabbit genome sequencing and advantages of rabbit reproduction (high cell numbers and yield in blastocysts, relatively late

implantation at a time when gastrulation is already proceeding, detailed morphologic and molecular knowledge on gastrulation stages and a hemochorial placenta structured similarly to the human placenta) are the reasons for the preferred use of a laboratory rabbit as a model for studying early embryo development (for review see Fischer *et al.*, 2012). Results obtained mainly on mouse and rabbit models are presented in this review.

Receptors for biogenic monoamines

Biogenic monoamine receptors are either ion channels or G protein-coupled receptor proteins. Receptor channels (ligand-gated ion channels) are composed of several transmembrane subunits, and binding the ligand induces opening of the channel. G protein-coupled receptors are composed of seven transmembrane domains connected by cytoplasmic and extracellular loops. Binding the ligand results in activation of GTP-binding proteins (G proteins), which then activate various effectors, such as adenylyl cyclases, phospholipase C beta, and other proteins involved in signal transduction (for review see Krauss, 2014).

Most information on the expression of biogenic monoamine receptors during preimplantation development period has been obtained on mouse and rabbit models. Several catecholamine (adrenergic and dopamine) serotonin and histamine receptors were detected in mouse and rabbit ovulated oocytes and preimplantation embryos (see Table 1). Some biogenic monoamine receptors were detected in oocytes and preimplantation embryos of other species as well (Čikoš *et al.*, 2014; Amireault and Dubé, 2005).

Embryonic stem cells (ESCs), derived from the inner cell mass of a blastocyst, can serve as an experimental model for studying early development. Although ESCs retain the high developmental potency of the founder embryo cells, several differences in the gene expression between preimplantation embryos and ESCs cells have been identified (Tanaka *et al.*, 2002; Tang *et al.*, 2010). Expression of catecholamine receptors has been examined in several mouse embryonic stem cell lines, and, in contrast to mouse preimplantation embryos, transcripts of all types of dopamine and adrenergic receptors were detected in mouse ESCs (Lee *et al.*, 2006; Kim *et al.*, 2008; Layden *et al.*, 2010; Čikoš *et al.*, 2015).

Differences in the expression of some adrenergic receptors between the spontaneously differentiating ESCs and undifferentiated ESCs indicate a role of these receptors in the process of embryonic stem cell differentiation (Čikoš *et al.*, 2015). Interestingly, catecholamine receptor types that couple primarily to G proteins with opposite actions on adenylyl cyclase activity (beta adrenoceptors, DR1 and DR5 stimulate adenylyl cyclase activity, and alpha 2 adrenoceptors, DR2, DR3 and DR4 inhibit adenylyl cyclase activity) are expressed in mouse and rabbit preimplantation embryos and in the mouse embryonic stem cells, indicating a cross-talk between signaling pathways activated by these receptors.

The role of biogenic monoamines in early embryo development

Experimental data indicate that biogenic monoamines can (besides their well-known neuro-

transmitter function) play an important role in basic developmental processes, such as embryogenesis and morphogenesis (for review see Pendleton *et al.*, 1998; Herlenius and Lagercrantz, 2001), and several biogenic monoamines have been identified in reproductive fluids of various mammalian species. For instance, adrenaline and noradrenaline have been detected in rabbit and bovine oviductal fluid and their concentrations varied with the region of the oviduct and the stage of the estrous cycle (Khatchadourian *et al.*, 1987; Way *et al.*, 2001).

Although the physiological role of biogenic monoamine receptors in early („pre-neural“) embryogenesis is mostly unknown, experimental data indicate that these receptors are functional and can be activated by agonists. Studies in mouse embryonic stem cells demonstrated that activation of dopamine and adrenergic receptors can trigger various signaling pathways influencing DNA synthesis and proliferation (Lee *et al.*, 2006;

Table 1. Receptors of biogenic monoamines detected in mouse and rabbit ovulated oocytes and preimplantation embryos

Receptor	Species	Stage (mRNA)	Stage (protein)	Reference
H2R	mouse	Blast	Blast	Zhao <i>et al.</i> , 2000
5-HT _{1D}	mouse	Ooc, Zyg, 2-cell, 8-16-cell, Blast	not tested	Veselá <i>et al.</i> , 2003 Il'ková <i>et al.</i> , 2004
5-HT ₇	mouse	Ooc, Zyg, 2-cell, 4-cell	Ooc, 4-cell	Amireault & Dubé 2005
α1B-AR	rabbit	Ooc	not tested	Čikoš <i>et al.</i> , 2014
α2A-AR	rabbit	Ooc, Morul, Blast	not tested	Čikoš <i>et al.</i> , 2014
α2C-AR	mouse	Ooc, 8-16-cell, Blast	Ooc, 8-16-cell, Blast	Čikoš <i>et al.</i> , 2007
	rabbit	Ooc, Morul, Blast	not tested	Čikoš <i>et al.</i> , 2014
β1-AR	rabbit	Ooc, Morul, Blast	not tested	Čikoš <i>et al.</i> , 2014
β2-AR	mouse	Ooc, 4-cell, 8-16-cell, Blast	Ooc, 4-cell, 8-16-cell, Bl	Čikoš <i>et al.</i> , 2014
	rabbit	Ooc, Morul, Blast	not tested	Chen <i>et al.</i> , 2011 Čikoš <i>et al.</i> , 2014
β3-AR	mouse	Ooc, 4-cell, 8-16-cell, Blast	not tested	Čikoš <i>et al.</i> , 2005
DR1	mouse	Blast	not tested	Čikoš <i>et al.</i> , 2015
DR2	mouse	Ooc, 8-16-cell, Blast	not tested	Čikoš <i>et al.</i> , 2015
DR3	mouse	Ooc, 4-cell, 8-16-cell, Blast	not tested	Čikoš <i>et al.</i> , 2015
DR4	mouse	4-cell, 8-16-cell, Blast	not tested	Čikoš <i>et al.</i> , 2015
DR5	mouse	Ooc, 4-cell, Blast	not tested	Čikoš <i>et al.</i> , 2015

Receptors: H2R, histamine receptor subtype 2; 5-HT_{1D}, serotonin receptor subtype 1D; 5-HT₇, serotonin receptor subtype 7; 5-HT_{2A}, serotonin receptor subtype 2A; α1B-AR, adrenergic receptor subtype alpha 1B; α2A-AR, adrenergic receptor subtype alpha 2A; α2C-AR, adrenergic receptor subtype alpha 2C; β1-AR adrenergic receptor subtype beta 1; β2-AR, adrenergic receptor subtype beta 2; β3-AR adrenergic receptor subtype beta 3; DR1-5, dopamine receptor subtypes 1-5.

Developmental stages: Ooc, unfertilized oocytes (metaphase II stage); Zyg, fertilized oocytes (zygotes); 2-cell, two-cell embryos; 4-cell, four-cell embryos; 8-16-cell, eight- to sixteen- cell embryos; Morul, morulas; Blast, blastocysts.

Kim *et al.*, 2008; Sun *et al.*, 2015). Most catecholamine receptors can regulate the intracellular cyclic adenosine monophosphate (cAMP) level, and it has been demonstrated that the cAMP signaling pathway can contribute to the regulation of mouse embryonic stem cell selfrenewal and differentiation (Faherty *et al.*, 2007; Layden *et al.*, 2010). Results of *in vitro* and *in vivo* experiments showed that catecholamine and serotonin receptors can activate signaling pathways regulating cell proliferation and survival in mouse preimplantation embryos (Markova *et al.*, 1990; Čikoš *et al.*, 2005; 2007; Veselá *et al.*, 2003; Il'ková *et al.*, 2004). Moreover, there are data suggesting that catecholamine receptors (embryonic or uterine) can participate in the process of blastocyst implantation (Henriquez *et al.*, 2006; Chen *et al.*, 2011). The involvement of histamine receptor subtype 2 (regulating cAMP level) in the process of embryo implantation has been documented in the mouse model as well (Zhao *et al.*, 2000). Results of *in vivo* experiments obtained on the mouse model indicate that catecholamine receptors together with glucocorticoid receptors could mediate effects of maternal stress on early embryo (Burkuš *et al.*, 2013; 2015; Janštová *et al.*, 2017; Zheng *et al.*, 2016; Tan *et al.*, 2017), and that expression of catecholamine receptors in early embryos can be influenced by maternal physiological status (Seeling *et al.*, 2017).

CONCLUSION

Mammalian preimplantation embryo is equipped with a variety of cell receptors indicating active communication between the early embryo and its environment. Results obtained in mouse and rabbit models indicate that except of receptors binding protein ligands, the receptors that bind small-molecule ligands, such as biogenic monoamines, are expressed in preimplantation embryos and in the embryonic stem cells derived from mouse blastocysts. Catecholamine receptors of all types can be expressed in mammalian preimplantation embryos, with some species-specific differences, and activation of these receptors can significantly influence preimplantation embryo development. Although the physiological role of biogenic monoamine receptors in early embryonic cells is not fully understood, experimental data

indicate their involvement in the regulation of cell proliferation, differentiation and survival under physiological as well as unfavorable or pathological conditions (e.g. during maternal stress).

ACKNOWLEDGEMENTS

This study was supported by the project No. ITMS 26220220204 of the Research & Development Operational Programme funded by the ERDF.

REFERENCES

- Amireault, P. & Dubé, F. 2005. Intracellular cAMP and calcium signaling by serotonin in mouse cumulus-oocyte complexes. *Molecular Pharmacology*, 68, 1678–1687.
- Babeřová, J., Šefčíková, Z., Čikoš, Š., Špirková, A., Kovaříková, V., Koppel, J., Makarevich, A. V., Chrenek, P. & Fabian, D. 2017. Exposure to neonicotinoid insecticides induces embryotoxicity in mice and rabbits. *Toxicology*, 392, 71–80.
- Burkuš, J., Čikoš, Š., Fabian, D., Kubandová, J., Czikková, S. & Koppel J. 2013. Maternal restraint stress negatively influences growth capacity of preimplantation mouse embryos. *General Physiology and Biophysics*, 32(1), 129–137.
- Burkuš, J., Kačmarová, M., Kubandová, J., Kokošová, N., Fabianová, K., Fabian, D., Koppel, J. & Čikoš, Š. 2015. Stress exposure during the preimplantation period affects blastocyst lineages and offspring development. *Journal of Reproduction and Development*, 61, 325–31.
- Chen, Q., Zhang, Y., Peng, H., Lei, L., Kuang, H., Zhang, L., Ning, L., Cao, Y. & Duan, E. 2011. Transient b2-adrenoceptor activation confers pregnancy loss by disrupting embryo spacing at implantation. *Journal of Biological Chemistry*, 286, 4349–4356.
- Čikoš, Š., Veselá, J., Il'ková, G., Reháč, P., Czikková, S. & Koppel, J. 2005. Expression of beta adrenergic receptors in mouse oocytes and preimplantation embryos. *Molecular Reproduction and Development*, 71, 145–153.
- Čikoš, Š., Reháč, P., Czikková, S., Veselá, J. & Koppel, J. 2007. Expression of adrenergic receptors in mouse preimplantation embryos and ovulated oocytes. *Reproduction*, 133, 1139–1147.
- Čikoš, Š., Czikková, S., Chrenek, P., Makarevich, A. V., Burkuš, J., Janštová, Ž., Fabian, D. & Koppel J. 2014. Expression of adrenergic receptors in bovine

- and rabbit oocytes and preimplantation embryos. *Reproduction in Domestic Animals*, 19, 92–100.
- Čikoš, Š., Fabian, D., Burkuš, J., Janštová, Ž. & Koppel, J. 2015. Expression of dopamine and adrenergic receptors in mouse embryonic stem cells and preimplantation embryos. *Biologia*, 70, 1263–1271.
- Diskin, M. G. & Morfia, D. G. 2008. Embryonic and Early Foetal Losses in Cattle and Other Ruminants. *Reproduction in Domestic Animals*, 43(Suppl.), 260–267.
- Fabian, D., Bystriansky, J., Čikoš, Š., Bukovská, A., Burkuš, J. & Koppel, J. 2010. The effect on preimplantation embryo development of non-specific inflammation localized outside the reproductive tract. *Theriogenology*, 74, 1652–60.
- Faherty, S., Fitzgerald, A., Keohan, M. & Quinlan, L. R. 2007. Selfrenewal and differentiation of mouse embryonic stem cells as measured by Oct4 expression: the role of the cAMP/PAK pathway. *In Vitro Cellular & Developmental Biology – Animal*, 43, 37–47.
- Fischer, B., Chavatte-Palmer, P., Viebahn, C., Navarrete Santos, A. & Duranthon, V. 2012. Rabbit as a reproductive model for human health. *Reproduction*, 144, 1–10.
- Fleming, T. P., Wilkins, A., Mears, A., Miller, D. J., Thomas, F., Ghassemifar, M. R., Fesenko, I., Sheth, B., Kwong, W. Y. & Eckert, J. J. 2004. Society for Reproductive Biology Founders' Lecture 2003. The making of an embryo: short-term goals and long-term implications. *Reproduction and Fertility*, 16, 325–337.
- Hardy, K. & Spanos, S. 2002. Growth factor expression and function in the human and mouse preimplantation embryo. *Journal of Endocrinology*, 172, 221–236.
- Henriquez, S., Tapia, A., Quezada, M., Vargas, M., Cardenas, H., Rios, M., Salvatierra, A. M., Croxatto, H., Orihuela, P., Zegers-Hochschild, F., Munroe, D. J. & Velasquez, L. 2006. Deficient expression of monoamine oxidase A in the endometrium is associated with implantation failure in women participating as recipients in oocyte donation. *Molecular Human Reproduction*, 12, 749–754.
- Herlenius, E. & Lagercrantz, H. 2001. Neurotransmitters and neuromodulators during early human development. *Early Human Development*, 65, 21–37.
- Humblot, P. 2001. Use of pregnancy specific proteins and progesterone assays to monitor pregnancy and determine the timing, frequencies and sources of embryonic mortality in ruminants. *Theriogenology*, 56, 1417–1433.
- Iľková, G., Reháč, P., Veselá, J., Čikoš, Š., Fabian, D., Czikková, S. & Koppel, J. 2004. Serotonin localization and its functional significance during mouse preimplantation embryo development. *Zygote*, 12, 205–213.
- Janštová, Ž., Burkuš, J., Kubandová, J., Fabian, D., Koppel, J. & Čikoš, Š. 2017. The effect of maternal stress on blastocyst quality depends on maternal physiological status. *General Physiology and Biophysics*, 36, 53–63.
- Khatchadourian, C., Menezo, Y., Gerard, M. & Thibault, C. 1987. Catecholamines within the rabbit oviduct at fertilization time. *Human Reproduction*, 2, 1–5.
- Khosla, S., Dean, W., Reik, W. & Feil, R. 2001. Culture of preimplantation embryos and its long-term effects on gene expression and phenotype. *Human Reproduction Update*, 7, 419–427.
- Kim, M. O., Na, S. I., Lee, M. Y., Heo, J. S. & Han, H. J. 2008. Epinephrine increases DNA synthesis via ERK1/2s through cAMP, Ca²⁺/PKC, and PI3K/Akt signaling pathways in mouse embryonic stem cells. *Journal of Cellular Biochemistry*, 104, 1407–1420.
- Krauss, G. 2014. *Biochemistry of Signal Transduction and Regulation*, 5th edn. Weinheim, Germany: Wiley-VCH Verlag GmbH, 2014.
- Kubandová, J., Čikoš, Š., Burkuš, J., Czikková, S., Koppel, J. & Fabian, D. 2014. Amount of maternal body fat significantly affected the quality of isolated mouse preimplantation embryos and slowed down their development. *Theriogenology*, 81, 187–195.
- Kwong, W. Y., Wild, A. E., Roberts, P., Willis, A. C. & Fleming, T. P. 2000. Maternal undernutrition during the preimplantation period of rat development causes blastocyst abnormalities and programming of postnatal hypertension. *Development*, 127, 4195–4202.
- Layden, B. T., Newman, M., Chen, F., Fisher, A. & Lowe, W. L. Jr. 2010. G protein coupled receptors in embryonic stem cells: a role for Gs- α signaling. *PLoS One*, 5, e9105.
- Lee, M. Y., Heo, J. S. & Han, H. J. 2006. Dopamine regulates cell cycle regulatory proteins via cAMP, Ca²⁺/PKC, MAPKs, and NF- κ B in mouse embryonic stem cells. *Journal of Cellular Physiology*, 208, 399–406.
- Markova, L. N., Sadykova, K. A. & Sakharova, N. I. U. 1990. The effect of biogenic monoamine antagonists on the development of preimplantation mouse embryos cultured *in vitro*. *Zhurnal evoliutsionnoĭ biokhimii i fiziologii*, 26, 726–732.
- Nepomnaschy, P. A., Welch, K. B., Mcconnell, D. S., Low, B. S., Strassmann, B. I. & England, B. G. 2006 Cortisol levels and very early pregnancy loss in humans. *Proceedings of the National Academy of Sciences of the United States of America*, 103, p. 3938–3942.
- Pendleton, R. G., Rasheed, A., Roychowdhury, R. & Hillman, R. 1998. A new role for catecholamines:

- ontogenesis. *Trends in Pharmacological Sciences*, 19, 248–251.
- Seeling, T., Čikoš, Š., Grybel, K. J., Janštová, Ž., Pendzialek, S. M., Schindler, M., Špirková, A. & Santos, A. N. 2018. A Diabetic Pregnancy Alters the Expression of Stress-Related Receptors in Gastrulating Rabbit Blastocyst and in the Reproductive Tract. *Reproductive Sciences*, 25, 174–184.
- Sun, F., Yang, X. J., Lv, H. Y., Tang, Y. B., An, S. M., Ding, X. P., Li, W. B., Teng, L., Shen, Y., Chen, H. Z. & Zhu L. 2015. β 2- Adrenoreceptor-mediated proliferation inhibition of embryonic pluripotent stem cells. *Journal of Cellular Physiology*, 230, 2640–2646.
- Tanaka, T. S., Kunath, T., Kimber, W. L., Jaradat, S. A., Stagg, C. A., Usuda, M., Yokota, T., Niwa, H., Rossant, J & Ko, M. S. 2002. Gene expression profiling of embryo-derived stem cells reveals candidate genes associated with pluripotency and lineage specificity. *Genome Research*, 12, 1921–1928.
- Tan, X. W., Ji, C. L., Zheng, L. L., Zhang, J., Yuan, H. J., Gong, S., Zhu, J. & Tan, J. H. 2017. Corticotrophin-releasing hormone and corticosterone impair development of preimplantation embryos by inducing oviductal cell apoptosis via activating the Fas system: an *in vitro* study. *Human Reproduction*, 32, 1583–1597.
- Tang, F., Barbacioru, C., Bao, S., Lee, C., Nordman, E., Wang, X., Lao, K. & Surani, M. A. 2010. Tracing the derivation of embryonic stem cells from the inner cell mass by single-cell RNASeq analysis. *Cell Stem Cell*, 6, 468–478.
- Thouas, G. A., Dominguez, F., Green, M. P., Vilella, F., Simon, C. & Gardner, D. K. 2015. Soluble ligands and their receptors in human embryo development and implantation. *Endocrine Reviews*, 36, 92–130.
- Ventura-Juncá, P., Irrázaval, I., Rolle, A. J., Gutiérrez, J. I., Moreno, R. D. & Santos, M. J. 2015. *In vitro* fertilization (IVF) in mammals: epigenetic and developmental alterations. Scientific and bioethical implications for IVF in humans. *Biological Research*, 48, 68.
- Veselá, J., Reháč, P., Mihalik, J., Czikková, S., Pokorný, J. & Koppel, J. 2003. Expression of serotonin receptors in mouse oocytes and preimplantation embryos. *Physiological Research*, 52, 223–228.
- Von Borell, E., Dobson, H. & Prunier, A. 2007. Stress, behaviour and reproductive performance in female cattle and pigs. *Hormones and Behaviour*, 52, 130–138.
- Way, A. L., Barbato, G. F. & Killian, G. J. 2001: Identification of norepinephrine in bovine oviductal fluid by high performance liquid chromatography. *Life Sciences*, 70, 567–576.
- Zhao, X., Ma, W., Das, S. K., Dey, S. & Paria, B. C. 2000. Blastocyst H2 receptor is the target for uterine histamine in implantation in the mouse. *Development*, 127, 2643–2651.
- Zheng, L. L., Tan, X. W., Cui, X. Z., Yuan, H. J., Li, H., Jiao, G. Z., Ji, C. L. & Tan, J. H. 2016. Preimplantation maternal stress impairs embryo development by inducing oviductal apoptosis with activation of the Fas system. *Molecular Human Reproduction*, 22, 778–790.